



## Tracheoesophageal fistula after intubation due to COVID-19: Endoscopic treatment

Enes Seyda Şahiner<sup>1</sup>, Bülent Ödemiş<sup>2</sup>, Tolga Canlı<sup>1</sup>

<sup>1</sup>Department Of Internal Medicine, Ankara City Hospital, Ankara, Turkey

<sup>2</sup>Department Of Gastroenterology, Ankara City Hospital, Ankara, Turkey

### ARTICLE INFO

Received Date:07.08.2022

Accepted Date:02.09.2022

### Keywords:

Tracheoesophageal fistula,  
COVID-19, OTSC

### ABSTRACT

Acquired non-malignant tracheoesophageal fistula (TEF) is a rare disease that has high levels of morbidity and mortality, and the etiology of TEF remains unclear. Better knowledge of predisposing factors such as traumatic intubation, highendotracheal tube cuff pressure, inflammation, and infection of COVID-19 could reduce the number of cases of acquired TEF. In this case report, a 36-year-old male patient who developed a TEF after intubation as a result of COVID-19 is presented. After short-term intubation, the TEF detected by neck CT and fluoroscopy was successfully closed with an endoscopic intervention using an over-the-scope clip and no leak was detected at the 6-month follow-up.

## Introduction

COVID-19 became an epidemic in the Chinese city of Wuhan in late 2019, often with atypical viral pneumonia. It may also present with a severe clinical condition that causes severe respiratory failure requiring intubation. The mechanical ventilation process is a difficult process to manage, which can cause many complications. TEF is one of the rare complications we encounter in this process.

As first described by Flege,<sup>1</sup> the most prevalent cause of tracheoesophageal fistula (TEF) is mechanical ventilation.<sup>2,3,4,5,6,7,8</sup>

Acquired TEF is a rare complication, developing and becoming symptomatic within 1-5 days after surgery, 5-15 days after trauma, 15-21 days after local infection, and 21-30 days after tracheal cuff-related injury.<sup>9</sup> TEF is rarely encountered upon short-term intubation in cases without trauma or surgery.<sup>10</sup>

Both cuff pressure and cuff volume (6-8 mL) should be monitored closely to minimize tracheoesophageal injuries in intubated patients. In this case report, we present a successful case of endoscopic TEF closure.

Written informed consent was obtained from the patient for publication of this text and accompanying images.

## Case Presentation

A 36-year-old male patient who was admitted to the service with COVID-19 PCR positivity and low oxygen saturation on 15.08.2021 developed acute respiratory failure due to COVID-19 pneumonia (Figure 1).

After 10 days of intubation, the patient was extubated and taken to the service from the intensive care unit. Recurrent aspiration attacks that particularly occurred after meals were noted in his follow-up appointments. The results of oral contrast-enhanced neck tomography were suspicious for TEF (Figure 2).




Figure 2. Neck CT 16.09.2021

Post Treatment Neck CT 27.09.2021



Figure 1. Torax CT Image (15.08.2021)

Fluoroscopy was therefore performed and TEF was detected (Figure 3).




Figure 3. Fluoroscopy 16.09.2021

Post Treatment Fluoroscopy 29.09.2021



Figure 4. Endoscopy 22.09.2021(11mm. OTSC)

An operation of approximately 30 minutes was formed on the patient by the endoscopist using an over-the-scope clip (OTSC) (Figure 4).

Subsequently, when the patient was checked by fluoroscopy and neck CT, no leakage was observed (Figure 2-3). He was transitioned to oral feeding and aspiration was not observed. The patient had no complaints at the 6-month follow-up visit.

### ***Discussion***

TEFs are epithelialized canals between the trachea and the esophagus. The causes of non-malignant acquired TEF include trauma, granulomatous infection, inflammation, previous trachea or esophageal operation, HIV infection,<sup>12,13,14</sup> poisoning and inhalation burns, tracheal tubes and intubation, and percutaneous tracheostomy.<sup>15</sup> In addition, intense inflammation caused by SARS-CoV-2 causes weakness in the tracheal wall structure because of mucosa and submucosa damage (16). Mucosal necrosis due to inflammatory infiltrates may result in a fistula between the tracheal tube cuff and the nasogastric tube.

Excessive movement of endotracheal tubes, diabetes, hypotension, malnutrition, steroid therapy, and long-term mechanical ventilation can be counted among the risk factors. The incidence of TEF has decreased, however, due to the advent of high-volume low-pressure cuffs.<sup>17,18</sup>

Many difficulties are encountered in the intraoperative and postoperative treatment of TEF in ENT and thoracic surgery clinics, and these cases come with possible complications along with high costs. There are studies showing the success of the use of OTSCs after failed stents and after operations in long-term follow-up.<sup>11</sup>

TEF is a situation in which many difficulties are encountered by otolaryngologists, gastroenterologists, and thoracic surgeons in the process of diagnosis and treatment. The use of OTSCs is a prominent and very comfortable method for treating TEF. It is predicted that OTSCs will be used more widely in the future for closure of perforations and fistulas that occur in the gastrointestinal system.<sup>7,8</sup>

In the management of TEF cases, there is a great lack of experience about the anatomy of the airway and digestive system, the size of the fistulas, and stent characteristics and types. When the dual stenting procedure is selected, both experience and knowledge are very important when combining an esophageal stent with an airway stent.

We hope that this case report will contribute to the literature on risk factors, clinical symptoms, diagnostic approaches, treatment methods, and prognosis in the context of TEF diagnosis and treatment.

### ***Declaration***

The authors have no conflicts of interest to declare.

## Reference

- 1- Flege Jr., J.B., Tracheoesophageal fistula caused by cuffed tracheostomy tube. *Ann Surg.* 1967;166:153-156
- 2- Bartlett R.H., A procedure for management of acquired tracheoesophageal fistula in ventilator patients. *J Thorac Cardiovasc Surg.* 1976;71:89-95
- 3- Hilgenberg A.D., Grillo H.C. Acquired nonmalignant tracheoesophageal fistula. *J Thorac Cardiovasc Surg.* 1983;85:492-498
- 4- Macchiarini P., Verhoye J.P., Chapelier A. et al. Evaluation and outcome of different surgical techniques for postintubation tracheoesophageal fistulas. *J Thorac Cardiovasc Surg.* 2000;119:268-276
- 5- Mathisen D.J., Grillo H.C., Wain J.C. et al. Management of acquired nonmalignant tracheoesophageal fistula. *Ann Thorac Surg.* 1991;52:759-765
- 6- Thomas A.N. et al. The diagnosis and treatment of tracheoesophageal fistula caused by cuffed tracheal tubes. *J Thorac Cardiovasc Surg.* 1973;65:612-619
- 7- Weiland T, Fehlker M, Gottwald T et al. Performance of the OTSC System in the endoscopic closure of gastrointestinal fistulae – a meta-analysis. *Minim Invasive Ther Allied Technol* 2012; 21: 249–258
- 8- Weiland T, Fehlker M, Gottwald T et al. Performance of the OTSC System in the endoscopic closure of iatrogenic gastrointestinal perforations: a systematic review. *Surg Endosc* 2013; 27: 2258–2274
- 9- Kaur D, Anand S, Sharma P, et al. Early presentation of postintubation tracheoesophageal fistula: Perioperative anesthetic management. *J Anaesthesiol Clin Pharmacol* 2012;28:114-6.
- 10- Rana R, Sapkota R, Shakya B, et al. Prolonged Intubation Induced Tracheoesophageal Fistula in Suspected Meningococcal Sepsis with ARDS: A Case Report. *JNMA J Nepal Med Assoc* 2018;56:980-982.
- 11- Nonthalee Pausawasdi et al., Successful Closure of a Benign Refractory Tracheoesophageal Fistula Using an Over-the-Scope Clip after Failed Esophageal Stent Placement and Surgical Management, *Clin Endosc* 2020;53:361-365 <https://doi.org/10.5946/ce.2019.106> Print ISSN 2234-2400 • On-line ISSN 2234-2443
- 12- Reed MF, Mathisen DJ. Tracheoesophageal fistula. *Chest Surg Clin N Am* 2003; 13: 271–89
- 13- Grillo HC. Surgery of the Trachea and Bronchi. Acquired Tracheoesophageal and Bronchoesophageal Fistula. London, BC Decker Inc. 2004;341–56
- 14- Ladurner R, Schulz C, Jacob P, Kuper M, Kratt T, Preyer S, Konigsrainer A. Surgical management of an esophagotracheal fistula as a severe late complication of repeated endoscopic stenting treatment. *Endoscopy* 2007; 39: E341–2.
- 15- Trottier SJ, Hazard PB, Sakabu SA, Levine JH, Troop BR, Thompson JA, McNary R. Posterior tracheal wall perforation during percutaneous dilational tracheostomy: an investigation into its mechanism and prevention. *Chest* 1999; 115: 1383–9
- 16- S. K. Mohanty, A. Satapathy, M. M. Naidu et al., “Severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) and coronavirus disease 19 (COVID-19) – anatomic pathology perspective on current knowledge,” *Diagnostic Pathology*, vol. 15, no. 1, p. 103, 2020.
- 17- Klainer AS, Turndorf H, Maewal H, Allender P. Surface alterations due to endotracheal intubation. *Am J Med.* 1975;58(5):674-83.
- 18- Stauffer JL, Olson DE, Petty TL. Complications and consequences of endotracheal intubation and tracheotomy. A prospective study of 150 critically ill adult patients. *Am J Med.* 1981;70(1)