



Hyperprolactinemia and galactorrhoea with duloxetine in neuropathic pain management

Nöropatik ağrı tedavisinde duloksetin ile ortaya çıkan hiperprolaktinemi ve galaktore

Eda DERLE, Ufuk CAN

Summary

Duloxetine is a serotonin-norepinephrine reuptake inhibitor that is widely used in chronic pain treatment in various disease. Hyperprolactinemia and galactorrhoea are rare side effects of this medication. Here, we reported a 34 year old female with multiple sclerosis who used duloxetine for pain management and mood disorder and experienced galactorrhoea.

Keywords: Duloxetine; galactorrhoea; hyperprolactinemia; mood disorder; neuropathic pain.

Özet

Duloksetin çeşitli hastalıklarda kronik ağrı tedavisinde sık olarak kullanılan serotonin-noradrenalin geri alım inhibitörüdür. Hiperprolaktinemi ve galaktore bu tedavinin nadir görülen bir yan etkisidir. Burada, multipl skleroz tanısı olan 34 yaşında bir kadın hastada ağrı ve duyu durumu bozukluğu nedeni ile duloksetin kullanımı ve medikal tedaviye bağlı gelişen galaktore olgusunu sunmaktayız.

Anahtar sözcükler: Duloksetin; galaktore; hiperprolaktinemi; duygudurum bozukluğu; nöropatik ağrı.

Introduction

Hyperprolactinemia is one of the most common cause of acquired hypogonadotropic hypogonadism and defined as elevated serum prolactin levels above the upper limits of normal range, with a prevalence of 1–1.5%.^[1, 2] Galactorrea was seen alone in approximately 30% and with amenorrhea in 75% of patient.^[1]

Hyperprolactinemia may be seen in the course of systemic disorders such as renal failure, hypothroidism, hypothalamic disease, pituitary disease and also commonly with drugs especially antipsychotic agents.^[1] The other drugs such as selective serotonin reuptake inhibitors and serotonin and noradrenalin reuptake inhibitors may cause the elevation of prolactin levels but it is not common.^[1, 2] The exact mechanism underlying this condition was not fully elucidated but in neuroendocrine system prolactin, dopamine, serotonin and noradrenaline interacts in several ways.^[1–3]

Duloxetine is a serotonin-norepinephrine reuptake inhibitor that is widely used in chronic pain treatment in various disease. Hyperprolactinemia and galactorrhoea are rare side effects of this medication, reported only in few case reports.

Case Report

Our patient was 34 years-old female, diagnosed with multiple sclerosis, on interferon beta 1b and gabapentin treatment. She used duloxetine (60 mg/d) for neuropathic and muscle pain and concomitant depressive symptoms for 3 months. In routine control she complained of milk like breast discharge that continued for one month. We considered this discharge as galactorrea and measured the serum prolactin level. The serum prolactin level was 93 ng/ml (Normal limits; 5.18–26.53 ng/ml). First we changed the duloxetine to escitalopram (10 mg/d) to eliminate the drug effect. Escitalopram also can rarely cause hyperprolactinemia, but we did not

Department of Neurology, Başkent University Faculty of Medicine, Ankara, Turkey

Submitted: 21.11.2018 Accepted after revision: 27.07.2019 Available online date: 04.05.2020

Correspondence: Dr. Eda Derle. Baskent Üniversitesi Tıp Fakültesi, Nöroloji Anabilim Dalı, Mareşal Fevzi Çakmak Caddesi, 10. Sokak, No: 45, Bahçelievler, Ankara, Turkey.

Phone: +90 - 312 - 203 68 68 / 1422 **e-mail:** edaderle@hotmail.com

© 2020 Turkish Society of Algology

Table 1. Review of case reports of duloxetine treatment and galactorrhea in literature

Case report	Dosage	Patient demographics	Indication	Symptom	Medical history	Evaluation	Prolactine level	Management	Outcome
Korkmaz et al. ^[8]	-	44 F	Depression	Milk-like breast discharge, pain and tenderness	Past usage of various antidepressant treatment	Normal pituitary gland magnetic resonance imaging Normal metabolic and biochemical blood tests	38.8 ng/ml	Switch to essitolapram and initiation of cabergolin 0.5 mg/twice a week	At the end of first month prolactin level was normal then cabergolin treatment stopped
Belli et al. ^[9]	60 mg	46 F	Depression	Milk-like breast discharge, pain, discomforting feeling of limbs at rest	-	Normal pituitary gland magnetic resonance imaging Normal metabolic and biochemical blood tests	37.9 ng/mL.	Switch to bupropion	Galactorrhea and restless legs symptoms improved after 2 weeks and normal prolactin level after six week
Demirci et al. ^[6]	30 mg 60 mg	28 F	Migraine	Milk like breast discharged	-	Normal pituitary gland magnetic resonance imaging Normal metabolic and biochemical blood tests	8.05 ng/mL (N)	Cessation of duloxetine	Galactorrhea stoped on the third day
Ashton et al. ^[10]		40 F	Dysthymic disorder	Milk like breast discharge	Past usage of various antidepressant galactorrhea with venlafaxin	Normal pituitary gland magnetic resonance imaging Normal metabolic and biochemical blood tests	28.2 ng/ml	Switch to bupropion and modofinil	Breast discharge discontinued and prolactin level decreased to normal level

prefer to cease all antidepressant treatment because of additional depressive disorder. Galactorrhoea stopped within days after cessation of duloxetine and also prolactin level decreased (39.78 ng/l) without additional treatment. At the same time we started to investigate endocrinologic causes of hyperprolactinemia. Metabolic tests and pituitary gland magnetic resonance imaging revealed normal results. Cabergolin treatment was initiated (0.25 mg twice in a week) by endocrinologist and the dosage was decreased in 3 months interval and prolactin level stayed within normal range. We thought that possible cause of the hyperprolactinemia in our case was antidepressant treatment.

Discussion

Duloxetine is widely used in neurology practice with diagnose of various pain disorders such as neuropathic pain, headache, fibromyalgia.^[4,5] Hyperprolactinemia and galactorrhoea are very rare side effects of these drugs (Table 1). Galactorrhoea may occur in even with normal prolactin level.^[6]

In a French pharmacovigilance cohort, in 11.863 patients who used serotonin reuptake inhibitors, hyperprolactinemia was reported as a side effect in 187 (1.6%).^[2] In this cohort fluvoxamine, citalopram, fluoxetine and paroxetine had higher ratios among them. Although there were several reports, they mentioned that duloxetine was not associated with an increased risk of hyperprolactinemia (1/225).^[2]

Antipsychotics are the most known drugs that are related to elevated prolactin levels and in the course of treatment, reaching up to 70% of the patients.^[7] Antipsychotic drugs showed their effect through dopamine receptor blockage. Dopamine is the main inhibitory factor for prolactin and this blockage causes elevation of the hormone levels.^[7] How do serotonin reuptake inhibitors cause hyperprolactinemia? It is known that serotonin is related to prolactin release.^[3] Several mechanisms proposed for this relation; one is via central 5-HT_{1C/2} receptors and the other is stimulation of GABAergic neurons situated near the tuberoinfundibular dopamine cells which have 5-HT_{1A} receptors on, and may increase prolactin levels through inhibition of inhibitory effect of dopamine in prolactin metabolism.^[3]

In drug induced hyperprolactinemia serum prolactin levels are usually below 100 ng/mL. The prolactin levels decrease in a few days after discontinuation of the related drug.^[1] If hyperprolactinemia persists or drug discontinuation is not appropriate, then the pituitary imaging can be performed to rule out adenoma.^[1] In symptomatic patients dose reduction or switch to another drug is recommended, if it's not possible, then low dose dopamine agonist is added to current treatment.^[1] Asymptomatic and mild elevation (<50 ng/ml) of prolactin may be followed without any treatment alteration.^[1]

In conclusion, while using these drugs we have to consider hyperprolactinemia in patients with any of the symptoms such as menstrual dysregulation, sexual dysfunction and breast discharge. Routine follow up of prolactin level is not recommended but in our opinion the symptoms may be reviewed in routine controls not to overlook this condition.

Informed Consent: *Written informed consent was obtained from the patient who participated in this study.*

Conflict-of-interest issues regarding the authorship or article: *None declared.*

Peer-review: *Externally peer-reviewed.*

References

1. Vilar L, Vilar CF, Lyra R, Freitas MDC. Pitfalls in the Diagnostic Evaluation of Hyperprolactinemia. *Neuroendocrinology*. 2019 Mar 20. Epub ahead of print
2. Trenque T, Herlem E, Auriche P, Drame M. Serotonin reuptake inhibitors and hyperprolactinaemia: a case/non-case study in the French pharmacovigilance database. *Drug Saf*. 2011 Dec 1;34(12):1161-6.
3. Coker F, Taylor D. Antidepressant-induced hyperprolactinaemia: incidence, mechanisms and management. *CNS Drugs*. 2010 Jul;24(7):563-74.
4. Curone M, Tullo V, Savino M, Proietti-Cecchini A, Bussone G, D'Amico D. Outcome of patients with chronic migraine with medication overuse and depression after duloxetine: influence of coexisting obsessive compulsive disorder. *Neurol Sci*. 2013 May;34 Suppl 1:S175-7.
5. Ferraro D, Plantone D, Morselli F, Dallari G, Simone AM, Vitetta F, Sola P, Primiano G, Nociti V, Pardini M, Mirabella M, Vollono C. Systematic assessment and characterization of chronic pain in multiple sclerosis patients. *Neurol Sci*. 2018 Mar;39(3):445-453
6. Demirci S, Unubol M, Demirci K. Galactorrhea with normal prolactin levels associated with duloxetine. *J Clin Psychopharmacol*. 2015 Jun;35(3):346-7.
7. Grigg J, Worsley R, Thew C, Gurchich C, Thomas N, Kulkarni

- J. Antipsychotic-induced hyperprolactinemia: synthesis of world-wide guidelines and integrated recommendations for assessment, management and future research. *Psychopharmacology (Berl)*. 2017 Nov;234(22):3279-3297.
8. Korkmaz S, Kuloglu M, Isik U, Saglam S, Atmaca M. Galactorrhea during duloxetine treatment: a case report. *Turk Psikiyatri Derg*. 2011 Fall;22(3):200-1.
 9. Belli H, Akbudak M, Ural C. Duloxetine-related galactorrhea and restless legs syndrome: a case report. *Psychiatr Danub*. 2013 Sep;25(3):266-7.
 10. Ashton AK, Longdon MC. Hyperprolactinemia and galactorrhea induced by serotonin and norepinephrine reuptake inhibiting antidepressants. *Am J Psychiatry*. 2007 Jul;164(7):1121-2.