

Broca's aphasia as a consequence of electric shock – A case study

Noorain Alam^{1,*} and Raj Kumar²

¹Post Graduate Institute Medical Education and Research, Chandigarh, India

²CU Shah Institute of Audiology and Speech Therapy, Surendranagar, Gujarat, India

ABSTRACT

The electric shock (ES) occurs upon contact of a human body part with any source of electricity that causes a sufficient current through the skin, muscles and hair. Broca's aphasia is a type of nonfluent aphasia, so called because speech production is halting and effortful. Current literature contains relatively little about the present state of knowledge concerning the neurological effects of electrical injury to man.

A 50 year old case reported C U Shah Medical College and Hospital, Surendranagar with complaint of electric shock in his right hand. He lost consciousness and when he regained consciousness he had right sided hemiparesis and had lost speech ability. The CT scan (fig 1) finding showed infarcts in left frontal lobe. The patient was assessed with the first part of the Western Aphasia Battery (WAB) (Kertesz, 1974). His spontaneous speech, repetition, naming are severely affected while auditory comprehension was relatively better. He was diagnosed with Broca's aphasia.

The pathology of Aphasia due to electrical shock involves blood vessel damages. Electrical shock can result in many neurological complications and Aphasias are one of them.

We need to assess all these possibilities while working with the persons who have suffered electrical shock.

Key Words: Electric shock, Broca's aphasia, speech therapy, Western Aphasia Battery test

Introduction

The electric shock (ES) occurs upon contact of a human body part with any source of electricity that causes a sufficient current through the skin, muscles and hair. Typically the expression is used to describe an injurious exposure to electricity. Electrical injury can result from direct contact with any source of electricity. While some electrical shocks may result in minor burns, there still may be serious internal damage hence, producing a complex pattern of injury and clinical manifestations. The exact pathophysiology of electrical injury is very difficult to predict due to the large number of variables that cannot be measured or controlled when an electrical current passes through tissue (1).

Accidental contact with exposed parts of electrical appliances or wiring, flashing of electric arcs from high-voltage power lines, lightning, machinery or occupational-related exposures or poking metal objects into an electrical outlet are all possible causes of electrical injury. High-voltage electrical trauma mainly observed in electrical workers produce some of the most devastating physical injuries. Repeated removal of the damaged tissue and extensive rehabilitation are

common while limb amputation rates for victims who experience direct electrical contact can be as high as 75%. In general, most victims who survive high-voltage electrical shock are left permanently disabled.

Away from the workplace, most electrical injuries are due to either indoor household low-voltage (less than 1000 V) electrical contact or outdoor lightning strikes. Domestic household 60-Hz electrical shocks are common and usually result in minor peripheral neurological symptoms or occasionally in skin surface burns. However, more complex injuries may result depending on the current path, particularly following biting or chewing on household appliance cord disclosures or outlets in small children. Compared to a high-voltage shock that usually is mediated by an arc, low-voltage shocks are more likely to produce a prolonged, "no-let-go" contact with the power source. This "no-let-go" phenomenon is caused by an involuntary, current-induced, muscle spasm.

The clinical effects on the nervous system resulting from electrical shock accidents can best be classified as immediate, secondary, and late manifestations (2). Immediate effects include loss of consciousness with confusion, headache, agitation, and convulsion. Effects without loss of

*Corresponding Author: Noorain Alam, M.D. Tutor, Speech & Hearing unit, ENT dept. Post Graduate Institute Medical Education & Research, Chandigarh, India, mobile: +91-9033 747078, E-mail: noorain.aslp@gmail.com

Received: 22.11.2013, Accepted: 08.05.2014

consciousness include deafness and severe pain. Motor signs including respiratory paralysis, tremor and complete paralysis. Sensory complaints include paresthesia and hypoaesthesia. Burns are also observed. Intermediate effects include temporary paralysis, pain in trunk, autonomic disturbances and headache. Late effects include effects on central as well as peripheral nervous system including aphasia, hemiplegia and cerebral dysfunction which were observed in our case. Electrical injury, also, often leads to problems with neurocognitive function, affecting speed of mental processing, attention, concentration, and memory. The high frequency of psychological problems is very well established and may be multifactorial in etiology (3). As with any traumatic and life-threatening experience, electrical injury may result in post traumatic psychiatric disorders which can be as life-changing as a major physical deformity (4). The larger currents can cause fibrillation of the heart and damage to the tissues and may lead to electrocution (5). Many people survive electrical trauma only to find a host of injuries including loss of consciousness, seizures, aphasia, visual disturbances, headaches, tinnitus, paresis, and memory disturbances (6).

Aphasia: Aphasia is a disturbance of the comprehension and formulation of language caused by dysfunction in specific brain regions (7).

Aetiology of Aphasia: It typically occurs after a stroke, head injury, brain tumour, infection or degenerative diseases leading to damage in areas of the brain cells controlling language. Stroke is the most common cause of aphasia, and it has been estimated that about 20-40% of stroke patients develop aphasia (8).

Classification of aphasia: Aphasics are generally grouped as being fluent or nonfluent (9). Nonfluent aphasics exhibit impaired flow of speech, effortful speech production with usually good comprehension skills, and varying degrees of difficulty with repetition. The nonfluent aphasics tend to display agrammatical or telegraphic speech (omission of function words) and depend primarily on the use of key content words (pain. . . leg). Nonfluent aphasia is typically seen in Broca's, transcortical motor, and global aphasia. Fluent aphasics exhibit a smooth flow of speech with varying degrees of deficits in comprehension and repetition. They may also exhibit circumlocution (speaking around a target word); paraphasic errors (use the word sister for mother or penkle for pencil); and neologisms (made up words e.g. 'gargot'). Typically Wernicke's, conduction,

transcortical sensory, and anomic aphasia are grouped in the fluent category.

Broca's Aphasia: It is a type of nonfluent aphasia, so called because speech production is halting and effortful. Damage is typically in the anterior portion of the left hemisphere. The dominant feature is agrammatism (impaired syntax). Content words (nouns, verbs) may be preserved but sentences are difficult to produce due to the problems with grammar, resulting in "telegraphic speech." In its more severe form, spoken utterances may be reduced to single words. Comprehension is typically only mildly to moderately impaired, and impairments are primarily due to difficulty understanding complex grammar. Repetition of words and sentences is usually poor.

In India there are very limited studies available on aphasia due to electric shock.

Case report

A 50 year old case was referred to the medicine department at C U Shah Medical College and Hospital, Surendranagar with complaint of electric shock in his right hand. He got electric shock when he was searching some household article due to the loose connection of electric switch board. He lost consciousness and when he regained consciousness he had right sided hemiparesis and had lost speech ability. The CT scan (Figure 1) finding showed infarcts in the left frontal lobe and he was diagnosed medicine CVA with hemiplegia. He was referred to speech therapy department for the loss of speech and language. The detailed speech and language evaluation was done.

Evaluation of speech and language: The patient was assessed with the first part of the Western Aphasia Battery (WAB) (Andrew Kertesz, 1974) (10). This part of WAB assesses fluency and information content of speech, naming, comprehension and repetition and writing. It is sufficient to determine the type of aphasia and it yields a measure of severity of aphasia, the Aphasia Quotient (AQ), ranging from 0 to 100, with scores above 93.8 being considered normal (non-aphasic). Scores obtained on each subtests are given in Table 1. Following were the findings:

Spontaneous speech: He was able to say only few words and needed phonemic and functional cues. He correctly responded to two items. He was expressing in single words which was effortful. Total score obtained was 5 out of 20.




Fig. 1. CT scan after electric shock.

Auditory verbal Comprehension: His auditory comprehension was reduced for both short and long phrases, and for yes/no questions. Auditory word recognition was also impaired. He was using few gestures and blinks for the items which he could not identify. Total scored obtained was 82 out of 200.

Repetition: His repetition was good for word level utterances but could not repeat sentence as

well complex sentence. Total obtained score was 14 out of 100.

Naming: In object naming he could name 3 items without any cue and 9 items with tactile and phonemic cues. In word fluency he could not name three animals in given one minute time while for the sentence completion task and responsive speech task he completed 3 items out of 5. Total score obtained was 29.

APHASIA QUOTIENT (AQ)–Total $\times 2 = 26.4$ suggestive of Aphasia (score < 93.8).

Total scores obtained on different subtests of WAB was calculated (table 2) and the patient was diagnosed with Broca’s aphasia.

Discussion

Current neurological literature contains relatively little about the present state of knowledge concerning the neurological effects of electrical injury to man. It is common knowledge that it is not the voltage which determines the effects of electrical shock on the human individual but rather the amount of current actually flowing through the body. In direct hand-to-hand current passages, fatalities are high (approximately 60%), whereas in hand-to-foot current passage fatalities are very much lower, around 20% (11). The pathology of Aphasia due to electrical shock involves blood vessel damages. Blood vessel

Table 1. WAB scoring on different subtests

SPONTANEOUS SPEECH	PATIENT SCORE	TOTAL SCORE
Information content	03	10
Fluency	02	10
Total	05	20
AUDITORY VERBAL COMPREHENSION		
Yes/no questions	46	60
Auditory word recognition	31	60
Sequential command	05	80
Total	82	200
REPETITION		
Total	14	100
NAMING		
Object naming	15	60
Word fluency	03	20
Sentence completion	05	10
Responsive speech	06	10
Total	29	100

Table 2. Total scores on WAB

FLUENCY	05
AUDITORY COMPREHENSION	4.1
REPETITION	1.4
NAMING	2.7

damage is fairly common and usually severe, especially involving large basal vessels; occasionally vessel rupture has been noted (12).

Pritchard (13) suggested that in the usual electrical accidents, if the victim is grounded, the current passes through the body producing heating and electrolytic and mechanical effects. However if grounding does not occur the electrical charge builds up in the body producing electrostatic effects. The electrostatic force upon a charged body is derived from mechanical repulsion that is exerted between all similarly charged particles and bodies. Thus all the separate constituents of the body that carry the charge will be repelled each other. The sudden expansile force of the body tissues acts as if the atmospheric pressure were suddenly reduced, giving rise to waves of decompression under the skin, and this is the force which affects the central nervous system tissues, most of the effects being evident where tissue rigidity or cohesion is least marked, as in the fluid spaces of the central nervous system. This may be possible pathology of Broca's Aphasia as post electrical shock injuries.

In conclusion, involvement of experienced consultants and a supportive environment are required to facilitate rehabilitation, return to gainful employment, and normal family and social functioning for the management of person with electrical shock injury. There are many neurological effects of electrical shock and aphasias are one of them. We need to assess all these possibilities while working with the person who have suffered electrical shock.

References

1. Mechanism of electrical injury. Chicago electrical trauma institute, <http://www.cetri.org/mechanism>. Accessed on November 15, 2013.
2. Silversides J. The neurological Sequelae of electrical injury. *Can Med Assoc J* 1964; 91: 195-204.
3. Grigorovich A, Gomez M, Leach L, Fish J. Impact of posttraumatic stress disorder and depression on neuropsychological functioning in electrical injury survivors. *J Burn Care Res* 2013; 34: 659-665.
4. Pliskin NH, Fink J, Malina A, et al. The neuropsychological effects of electrical injury. New insights. *Ann N Y Acad Sci* 1999; 888: 140-149.
5. Reilly Patrick J. Applied bioelectrical stimulation to electro pathology, 2nd edition. Springer. ISBN 1998; 978-0-387-98407-0.
6. Pliskin NH, Meyer GJ, Dolske MC, et al. Neuropsychiatric aspects of electrical injury. A review of neuropsychological research. *Ann N Y Acad Sci* 1994; 720: 219-223.
7. Damasio AR. Aphasia. *N Engl J Med* 1992; 326: 531-539.
8. Wade DT, Hewer RL, David RM, Enderby PM. Aphasia after stroke: natural history and associated deficits. *J Neurol Neurosurg Psychiatry* 1986; 49: 11-16.
9. Kearns KP. Broca's aphasia. In: La Pointe LL, editor. *Aphasia and related neurogenic language disorders*. 2nd edition. New York7 Thieme Medical Publishers; 1997; p. 1-34.
10. Kertesz A, Poole E. The aphasia quotient: the taxonomic approach to measurement of aphasic disability. *Can J Neurol Sci* 1974; 1: 7-16.
11. Boxall EA, Lauener RW, McIntosh HW. Atypical manifestations of hyperthyroidism. *Canad Med Ass J* 1964; 91: 204-211.
12. Hassin GB. General pathological considerations in brain injury, In: *Injuries to the brain and spinal cord and their coverings; neuropsychiatric, surgical, and medico-legal aspects*, 3rd ed., edited by S. Brock, Williams & Wilkins Company, Baltimore, 1949; p. 25.
13. Pritchard EAB. Changes in the central nervous system due to electrocution. *Lanset* 1934; 223: 1163-1167.