

DEMOGRAPHIC EVALUATION OF LARYNGO-TRACHEAL STENOSIS FOLLOWING PROLONGED ENDOTRACHEAL INTUBATION

HASSAN ABSHIRINI*
MOHAMMAD R. PIPELZADEH**
KHASHAYAR AHMADI*

SUMMARY: Prolonged trans-laryngeal intubation is associated with increased laryngeal injury, glottic and sub-glottic stenosis, infectious complications, and tracheal injury such as tracheomalacia, and tracheal stenosis. Several studies assessing the laryngeal complications are available in Caucasoid populations. Minimal data are available in the literature related to Iran. In this study we have evaluated the etiological factors of laryngo-tracheal stenosis following intubation.

It was a prospective analytical study in Otolaryngology- Head and Neck Surgery Department, Ahwaz Jondishapour University of Medical Sciences, from 2004 until 2006 (19 months) in Ahwaz, Iran. Ninety eight patients (61 patients were male) who were intubated for more than 48 hours in Imam Khomeini, Golestan and Razi University Hospitals critical care unit with different causes and were referred to otolaryngology emergency ward with respiratory distress (The prevalent cause was head trauma) were enrolled in the study. All patients underwent laryngeal video endoscopy and laryngeal computed tomography scan at by an otolaryngologist surgeon who was blinded to the intubation variables.

Twenty five patients had been presented by laryngo-tracheal stenosis and the most common site of that was sub-glottic area (sixteen patients). Statistical analytic findings showed that from several factors such as tube size ($P=0.088$), skill of intubating doctor (consultant or registrar; $P=0.146$), duration of intubation ($P=0.002$) and emergency intubation ($P=0.240$) only prolonged intubation ($P=0.002$) was associated with higher incidence of laryngo-tracheal stenosis.

In conclusion, laryngo-tracheal stenosis after prolonged endotracheal intubation was directly associated with duration of intubation. The most common site of laryngeal stenosis was sub-glottic area.

Keywords: Endotracheal intubation, laryngo-tracheal stenosis

INTRODUCTION

Prolonged trans-laryngeal intubation is associated with increased laryngeal injury, glottic and sub-glottic stenosis, infectious complications, and tracheal injury such as tracheomalacia, and tracheal stenosis (1,2). Converting intubation to a tracheotomy also may facilitate suctioning, feeding, and mobility; promote early return of speech; and

decrease the work of breathing (2). Patients with marginal ventilatory capacity by reducing the airway resistance (1,2). Meticulous surgical technique, aggressive treatment of postoperative infections, and the use of the high-volume, low-pressure cuffed tube help to minimize the risk of tracheal stenosis (3). Evidence suggests that the risk of long-term airway complications significantly increases with trans-laryngeal intubation beyond the tenth day (4,5). Tracheal stenosis and sub-glottic stenosis are complications predisposed by previous endotracheal tube intubation, high tracheostomy or cricothyroidotomy, and trauma to the

*From Department of Otorhinolaryngology, Head and Neck Surgery, Imam Khomeini Hospital, Ahwaz Jondishapour University of Medical Sciences, Ahwaz, Iran.


**From Department of Anaesthesiology, Imam Khomeini Hospital, Ahwaz Jondishapour University of Medical Sciences, Ahwaz, Iran.

airway. Patients at increased risk for tracheal stenosis include children and patients tracheotomized for closed head trauma. While the indications for intubation are varied, it is commonly used for ventilatory assistance, relief of upper airway obstruction and to aid in tracheo-bronchial toileting. With the advent of high volume low-pressure tubes, it is common for the patients to be ventilated through the endotracheal tube for up to three weeks. The complication of intubation depends on multiple factors and can also at times be a major source of morbidity to the patient during recovery. Laryngeal complications after prolonged intubations have varied incidence rates in the literature. Hoarseness of voice, cough, dysphagia, aspiration and stridor can develop after extubation, immediately. Laryngeal evaluation of such patients becomes necessary to assess the nature of airway injury. Common injuries following long-term intubation can manifest in the form of erythema, ulceration, granulomas (Figure 1), arytenoid dislocations, (1-3) sub-glottic stenosis and vocal fold immobility (6-12). While several studies assessing the laryngeal complications are available in Caucasoid populations; minimal data is available in the literature related to Iran. Therefore we evaluated the etiologic risk factors of laryngo-tracheal stenosis in patients after prolonged intubation to determine the association with risk factors. The medical ethics committee of the Otorhinolaryngology-Head and Neck Surgery Department approved the study and informed consent was obtained from all subjects. All references recommended that the conversion from trans-laryngeal intubation to tracheotomy be made as early as possible to minimize the duration of trans-laryngeal intubation (17,18).

MATERIALS AND METHODS

In this prospective analytical study 98 (sixty one patients were male) patients who were referred to ENT (Ear Nose Throat) emergency wards of Imam Khomeini, Golestan and Razi University Hospitals with respiratory distress were included. All ninety eight patients have age >20 years old, extubated at the intensive care unit of above named Hospitals after at least 48 hour of intubation were included in the study. Laryngeal examination with a rigid 70 degree and direct laryngeal video-endoscopy was done for all patients, by the ENT surgeon who was blinded to the intubation variables. Laryngeal computed tomography scan (CT scan) was done for all patients for evaluation of sub glottic stenosis. The association of laryngeal findings with demographic details along with intubation variables nature of intubation (emergency or elective), size of endotracheal tube, skill of intubating personnel (Consultant/ Resident), and cuff pressure during the intubated period, total period of intubation were studied.

Figure 1: Laryngeal sequelae distribution.



Statistical Analysis

The results were analyzed by using SPSS (version: 13) statistical software. Parametric (normally distributed) data were analyzed using *t*-test for group comparisons. Nonparametric (skewed) data were analyzed using Kruskal-Wallis test. Univariate correlation between presence of laryngeal complications and parametric variables was done using Spearman's coefficient of correlation. Chi-square test was used to analyze datum and p value of (<0.05) was accepted as being statistically significant.

RESULTS

Laryngeal abnormalities were seen in thirty six of the ninety eight patients during examination on post extubation period by ENT surgeon with helping laryngeal rigid endoscopy and laryngeal CT scan. At the end of laryngeal rigid endoscopy, physical examination and CT scan findings revealed that 25 patients had laryngo-tracheal stenosis. Vocal cord granuloma was the most common feature in the set of abnormal findings on laryngeal rigid endoscopy in six patients; other abnormalities were arytenoids dislocation in three patients and unilateral vocal cord palsy in two patients. All of individual factors such as age, sex, height, and weight had not any significant analytical relationship with laryngeal complication. Table 1 shows data regarding intubation variables and their association with the laryngeal sequelae on laryngeal endoscopy in patients. Multivariate and Univariate analysis of the intubation variables showed that between several factors include: larger endotracheal tube size ($P=0.088$), skill of intubating doctor (consultant or registrar; $P=0.146$), nature of intubation (Table 2) (elective or emergency; $P=0.240$) and prolonged intubation ($P=0.002$) (Table 3), the presence of laryngeal complications (laryngo-tracheal stenosis) associated with the laryngeal rigid endoscopy and CT scan findings after extubation was related only to the duration of intubation significant factors ($P=0.001$). The most

Table 1: Statistics intubation attributes.

Variable	P value
Intubating doctor	0.146
Nature of intubation	0.240
Tube size	0.088
Duration of intubation	0.002

Table 2: Intubation nature and laryngeal complication.

Nature of intubation	Laryngo-tracheal stenosis	Other complication
Elective	4	-
Emergency	8	5
Consultant	5	3
Resident	8	3

Table 3: Intubation duration and laryngeal complication .

Intubation duration	Laryngo-tracheal stenosis	Other complication
Less than 7 days	-	6
Between 7 and 14 days	6	3
More than 14 days	19	2

common site of laryngeal involvement was sub-glottic stenosis. Twenty five patients had tracheolaryngeal stenosis, with laryngeal CT scan and endoscopy findings. In this study the most common site of laryngeal stenosis was sub-glottic area (sixteen patients).

DISCUSSION

Laryngeal injuries following intubation have a reported incidence from 60% to 90% and permanent sequelae are reported to be about 15% in the world literature (5). The time to switch over from endotracheal tubes to tracheostomy is controversial (6). Current data suggest that ventilation through high volume low-pressure endotracheal tubes are the method of choice of artificial airway up to 3 weeks (11). The laryngeal injuries can vary from mucosal injuries like vocal cord erythema, edema, granulations and ulcerations to more permanent sequelae like vocal cord palsy (4-6), arytenoids dislocation (1-3), and sub-glottic stenosis (4-10). In our study vocal cords and the arytenoids were not the only components of the larynx involved in injuries. There were patients with vocal cord paralysis, arytenoids dislocations or tracheal and sub-glottic stenosis in this study population. Laryngeal injuries are seen in

patients intubated with larger size endotracheal tube. The determination of tube size of an adult patient is usually based on the physician's assessment of neck morphology and external features of the larynx. There is no standard formula to serve as a guide to appropriate tube size. A size eight endotracheal tube is generally used for adults and size 7 or 7.5 for small built adults. In this study the size of endo-tracheal intubation tube had not any significant statistical relation with tracheolaryngeal stenosis ($P=0.088$). Literature (10,14). shows increased incidence of laryngeal injuries with higher cuff pressures. In our study population, 90% of the patients had a standard cuff pressure of 30 cm H₂O. The cuff restricts the flow of blood through the tracheal tissues thus causing ischemic damage. The use of high-volume, low-pressure endotracheal tubes are also a routine practice in our intensive care unit. These tubes have a cuff with a large resting volume and diameter and a thin compliant wall that allows a seal with the trachea to be achieved without stretching its wall. This reduces the risk of injury to tracheal tissues; thereby reducing the risk of significant cuff-induced complications following prolonged intubation. The association of more laryngeal complications with emergency intubations is an expected occurrence.

Intubations done in the emergency sitting have the disadvantages of less preparation, inadequate patient sedation and less co-operation on the part of the patient that can be responsible for higher occurrence of laryngeal injuries. In this study the emergency intubating had not any significant statistical relation with tracheolaryngeal stenosis ($P=0.240$). The skill of the intubating physician also plays an important role in the genesis of laryngeal injuries. The residents and trainees with less skill and experience have comparatively greater association with complications than the consultants. Use of appropriate tube size and monitoring of duration of intubation is essential in the prevention of laryngeal complications ($P=0.002$). Only about 20% of the patients had permanent sequelae. This study shows that, at the post extubation period, duration of intubation was the only predictor of post extubation laryngeal sequelae. Use of appropriate tube size and monitoring of duration of intubation are of paramount importance. A systematic laryn-

geal examination after extubation, whenever indicated, decreases the secondary sequelae by initiating appropriate measures when necessary. It is highly suggested to early otorhinolaryngology consultation for the patients who are intubated in intensive care unit (ICU) for any etiologies. It seems to large analytical studies with bigger cohorts are needed to define the role of medical disorders and other parameters in producing laryngeal complications in patients with prolonged intubation.

ACKNOWLEDGMENTS

Authors acknowledge Dr Soltanzadeh, the president of Ahwaz Jondi-Shapour University of Medical Science for permitting the study to be conducted in all hospitals and for his suggestions, Dr. Mohammad Reza Ghoushe Consultants in Department of Critical Care, for permitting the study to be conducted in the ICU and also helping in the study design.

REFERENCES

1. Rudert H : Uncommon injuries of the larynx following intubation, Recurrent paralysis, torsion and luxation of the cricoarytenoid joints. *HNO*, 32:393-398, 1984.
2. Rieger A, Hass I, Gross M, Gramm HJ, Eyrich K : Intubation trauma of the larynx: A literature review with special reference to arytenoid cartilage dislocations. *Anesthesiol Intensivmed Notfallmed Schmerzther*, 31:281-287, 1996.
3. Obrebowski A, Wojnowski W : Hoarseness resulting from post-intubation arytenoid cartilage subluxation in an 11-year-old girl. *Otolaryngol Pol*, 52:223-226, 1998.
4. Lundy DS, Casiano RR, Shatz D, Reisberg M, Xue JW: Laryngeal injuries after long-term intubation. *J Voice*, 12:360-365, 1998.
5. Santos PM, Afrassiabi A, Weymuller EA Jr : Risk factors associated with prolonged intubation and laryngeal injury. *Otolaryngol Head Neck Surg*, 111:453-459, 1994.
6. Panda NK, Mann SB, Raja BA, Batra YK, Jindal SK : Fiberoptic assessment of post intubation laryngotracheal injuries. *Indian J Chest Dis Allied Sci*, 38:241-247, 1998.
7. Benjamin B : Prolonged intubation injuries of the larynx: Endoscopic diagnosis, classification and treatment. *Annal Otol Rhinol Laryngol*, 160:1-15, 1993.
8. Colice GL : Resolution of laryngeal injury following translaryngeal intubation. *Am Rev Respir Dis*, 145:361-364, 1992.
9. Rajagopalan R, Rajaram R, Pai M, Mahendran S, Praveen RJ, Baskaran A : Incidence and risk factors of post-extubation stridor in a cohort of adult intensive care patients in South India (abstract). *Indian J Crit Care Med*, 3:75-76, 1999.
10. Kastanos N, Estopa Micro R, Marin Perez A, Xaubet Mir A, Augusti-Vidal A: Laryngotracheal injury due to endotracheal intubation: Incidence, evolution and predisposing factors: A prospective long-term study. *Crit Care Med*, 11:362-367, 1983.
11. Hsu CL, Chen KY, Chang CH, Jerng JS, Yu CJ, Yang PC: Timing of tracheostomy as a determinant of weaning success in critically ill patients: A retrospective study *Crit Care*, 9:R46-52, 2005.
12. Ellis SF, Pollak AC, Hanson DG, Jiang JJ: Videolaryngoscopic evaluation of laryngeal intubation injury: Incidence and predictive factors. *Otolaryngol Head Neck Surg*, 114:729-731, 1996.
13. Volpi D, Lin PT, Kurloff DB, Kimmelman CP : Risk factors for intubation injury of the larynx. *Annal Otol Rhinol Laryngol*, 96:684-686, 1987.
14. Granja C, Faraldo S, Laguna P, Gois L : Control of the endotracheal cuff balloon pressure as a method of preventing laryngotracheal lesions in critically ill intubated patients. *Rev Esp Anesthesiol Reanim*, 49:137-140, 2002.
15. Buckels NJ, Khan ZH, Irwin ST and others: Post-operative sputum retention treated by minitracheostomy-a ward procedure. *Br J Clin Pract*, 44:169-171, 1990.
16. Heffner JE : The role of tracheotomy in weaning, *Chest*, 20:4778- 4815, 2001.
17. Valsh ME, Shorten GD : Preparing to perform an awake fiberoptic intubation. *Yale J Bioi Med*, 71:537-549, 1998.
18. Whited RE : A prospective study of laryngotracheal sequelae in long-term intubation. *Laryngoscope*, 94:367-377, 1994.

Correspondence:

Khashayar Ahmadi
Department of Otorhinolaryngology,
Head and Neck Surgery,
Imam Khomeini Hospital,
No. 50 Shohada Street,
Ahwaz, IRAN.
e-mail: dr.khashayar.ahmadi@gmail.com