

Effect of intra-abdominally administered mesalazine (5-aminosalicylic acid) in experimental peritonitis

Aziz Sener,¹ Alper Sahbaz,¹ Leyla Turker Sener,² Merva Soluk Tekkesin,³ Bulent Kaya⁴

¹Department of General Surgery, H.S.U Kanuni Sultan Suleyman Training and Research Hospital, Istanbul, Turkey

²Department of Biophysics, Istanbul University Faculty of Medicine, Istanbul, Turkey

³Department of Institute of Oncology, Istanbul University, Istanbul, Turkey

⁴Department of General Surgery, H.S.U Fatih Sultan Mehmet Training and Research Hospital, Istanbul, Turkey

ABSTRACT

OBJECTIVE: In this study, the effect of mesalazine was studied on TNF alpha, IL-1 beta, IL-6, and C-reactive protein (CRP) levels and inflammatory changes in rat lungs with experimental peritonitis.

METHODS: In total, 24 male Sprague–Dawley rats weighing 250–280 g were used in the study. The rats were divided into three groups based on no irrigation or irrigation with isotonic solution or mesalazine. Secondary peritonitis was generated by cecum penetrations. Group I, no irrigation was given after the development of peritonitis; Group II, irrigation was performed using isotonic solution 24 h after the development of peritonitis; Group III: irrigation was performed using mesalazine 24 h after the development of peritonitis. Blood samples were taken in the 48th hour for measuring TNF alpha, IL-1 beta, IL-6, and CRP levels. Lung tissue samples were taken for examining the effect of mesalazine in the development of systemic sepsis.

RESULTS: TNF alpha, IL-1 beta, and CRP levels were significantly low in Group III than in the other groups ($p < 0.005$). In the histologic examination, leucocyte infiltration in the lung was found low in Group III.

CONCLUSION: TNF alpha, IL-1 beta, and CRP levels and leucocyte infiltration in the lung were found to be low in rats that were administered peritoneal irrigation using mesalazine after the development of secondary peritonitis. Peritoneal irrigation using mesalazine may be useful in patients requiring surgery due to secondary peritonitis.

Keywords: Peritonitis; TNF alfa; IL-1 beta; IL-6; mesalazine.

Cite this article as: Sener A, Sahbaz A, Turker Sener L, Soluk Tekkesin M, Kaya B. Effect Of Intra-Abdominally Administered Mesalazine (5-Aminosalicylic Acid) In Experimental Peritonitis. *North Clin Istanbul* 2018;5(3):171–175.

Sepsis is a serious pathology with significant morbidity and mortality [1]. Some proinflammatory mediators indirectly cause sepsis as a response against exposure to microbial products. Cytokine response is associated with morbidity and mortality in sepsis [2, 3]. Microbial products include liposaccharides (endotoxins or lipopolysaccharides) produced by gram-negative microorganisms, peptidoglycans and teichoic acids produced by gram-positive microorganisms, cell-wall components and many others produced by fungi and yeasts.

Secondary peritonitis is a common morbidity that requires surgical intervention. Gastrointestinal system-related intra-abdominal contamination initiates the cytokine synthesis. Cytokines TNF α and IL-1 β lead the pathogenesis of sepsis. These cytokines increase in the first 4 h after the development of sepsis. They reach the highest level in the 24th hour. C-reactive protein (CRP) is an acute-phase reactant, and CRP levels increase in sepsis.

Mesalazine or 5-aminosalicylic acid (5-ASA) is an anti-inflammatory and antioxidant drug that has been

Received: August 15, 2017 *Accepted:* October 10, 2017 *Online:* June 08, 2018

Correspondence: Dr. Aziz SENER. H.S.U Kanuni Sultan Suleyman Egitim ve Arastirma Hastanesi, Genel Cerrahi Anabilim Dalı, Istanbul, Turkey

Phone: +90 532 436 05 53 e-mail: drazisener@hotmail.com

© Copyright 2018 by Istanbul Provincial Directorate of Health - Available online at www.northclinist.com



used in treatment of inflammatory bowel diseases for approximately 50 years [4, 5]. 5-ASA is a small hydrophilic organic acid and is well absorbed from the intestines. It is rapidly transformed to N-acetyl in the intestinal epithelium and liver, and the inactive molecule N-acetyl-5-ASA is generated. Some parts of the inactive metabolite are secreted back to the lumen and removed from the body by feces. Some parts of 5-ASA are metabolized in the liver and extracted by the kidney. However, the clinical effect of 5-ASA is not associated with systemic absorption and redistribution on target organs; its topical effect is observed in the colon. There are various views on the effect mechanisms of 5-ASA complexes. It was stated that 5-ASA complexes inhibit IL-1 and IL-2 synthesis in the inflammatory period. It was also proved that they inhibit TNF- α synthesis [6].

The aim of the present study was to investigate the effect of intra-abdominally administered mesalazine on cytokines (TNF- α , IL-1 β , and IL-6) and on the level of acute-phase reactant CRP during sepsis due to secondary peritonitis.

MATERIALS AND METHODS

Animal study

The study was conducted at the Institute of Experimental Medicine, Istanbul Faculty of Medicine, Istanbul University. The study protocol was approved by the Animal Care Ethics Committee. All procedures were conducted in accordance with the ethics guidelines for the treatment and welfare of experimental animals by the Istanbul Faculty of Medicine, Istanbul University, and Helsinki declaration. In total, 24 male Sprague–Dawley rats weighing 250–280 g were used in this study. The animals were housed at 21°C and were given tap water and standard rat food ad libitum.

Surgical procedure

The animals were anesthetized via an intramuscular injection of ketamine hydrochloride (50–100 mg/kg of body weight). Twenty-four male rats were randomly assigned to three groups. Group I, no irrigation after the development of peritonitis; Group II, irrigation using isotonic solution in the 24th hour after the development of peritonitis; and Group III, irrigation using mesalazine solution in the 24th hour after the development of peritonitis.

Peritonitis was generated by small holes on the cecums of rats using a 22-G needle after laparotomy. Ten




FIGURE 1. (A, B) Midline laparotomy (C) Peritoneal lavage with mesalazine (D) Closure of the abdominal cavity.

milliliter isotonic solution was abdominally administered to Group II and mesalazine (0.2 g/10mL) to Group III 24 h later (Fig. 1). A second laparotomy was performed in the 48th hour. Intracardiac blood samples were taken. Blood samples were collected for the measurement of TNF α , IL-1 β , IL6, and CRP levels. All rats were sacrificed 48 h after the first laparotomy. Tissue samples were taken from the lung.

Cytokines measurement

Samples were collected in serum separator tubes. After clot formation, samples were centrifuged at 1000 \times g for 10 min, and serum was collected. Serum was stored at -40°C until required for analysis. Cell signal proteins were assessed in the top supernatant of blood by TNF- α , IL-1 β , IL-6, and CRP levels using an enzyme-linked immunosorbent assay kit in accordance with the manufacturer's guidelines.

Histologic analysis

Lung tissue samples were taken from the rats for histopathologic investigation because the risk of development of multiple organ failure was mainly in the lung in sepsis. For histologic assessments, lung samples were obtained from eight anesthetized rats in each group at 48 h through excisional biopsy. All specimens were fixed in 10% buffered formalin. Paraffin blocks were prepared from routinely processed specimens, and 5- μ m sections were cut and deparaffinized. The sections were stained using hematoxylin–eosin (H&E). Histopathologic examinations were performed by a blinded pathologist with light microscopy at \times 200 magnification.

TABLE 1. Characteristics of groups

	N	Mean	Median	St. Dev.	Min.	Max.
TNF	18	227.4	227.3	27.8	185.1	175.4
IL-6	18	9.8	9.8	1.3	7.9	12.6
IL-1beta	18	1827.8	1767.5	196.9	1556.3	2271.0
CRP	18	184.2	198.1	31.3	128	237.4
Lenfosit	14	48.4	39.5	24.1	9.0	92.6
Nötrofil	13	19.4	11.4	21.7	1.5	63.5
Lökosit	14	6.9	7.2	2.6	3.0	12.3

TABLE 2. Post-hoc test

		p
TNF	1-2	0.073*
	1-3	0.003*
	2-3	0.038*
IL-1beta	1-2	0.214*
	1-3	0.419*
	2-3	0.914*
CRP	1-2	0.808*
	1-3	0.001*
	2-3	0.114*

*Mann-Whitney U p value.

Statistical method

Descriptive statistics were used to describe continuous variables (mean, standard deviation, minimum, maximum, and median). Comparison of three independent and non-normally distributed continuous variables was performed using the Kruskal–Wallis test. For significant results of the Kruskal–Wallis and Mann–Whitney U test, Bonferroni correction was used as a post-hoc analysis test.

Statistical significance level was set at 0.05. Statistical analyses were performed using MedCalc Software version 12.7.7 (MedCalc Software bvba, Ostend, Belgium; <http://www.medcalc.org>; 2013).

RESULTS

Intracardiac blood samples were taken 48 h after the operation for TNF- α , IL-1 β , IL-6, and CRP measurement. TNF- α , IL-1 β , and CRP levels were detected at a significantly lower level in Group III than in the other two

groups (Kruskal–Wallis, $p < 0.05$) (Table 1). According to the the post-hoc test results, the differences stemmed from Groups I and III (Mann–Whitney U, $p < 0.016$, Bonferroni correction) (Table 2).

There is significant difference between TNF and Group. (Kruskal-Wallis $p < 0.05$) According to the Post-Hoc test results, the differences stems from 1. and 3. groups. There is significant difference between IL-1beta and Group. (Kruskal-Wallis $p = 0.041$) According to the Post-Hoc test results, the differences are not caused by two groups. There is significant difference between CRP and Group. (Kruskal-Wallis $p = 0.011$) According to the Post-Hoc test results, the differences stems from 1. and 3. groups (Mann-Whitney U $p < 0.016$ Bonferroni correction) (Table 3).

The histologic sections were examined under light microscopy and assessed for inflammation in lung tissue, which can be an indirect sign of sepsis. Slight differences of inflammatory reactions were determined between the groups. Group I showed more inflammatory infiltration than Groups II and III. Group III showed relatively lesser inflammatory infiltration than Group II (Fig. 2).

DISCUSSION

Secondary peritonitis is one of the main causes of sepsis and multiple organ failure. Its mortality rate is still high [7]. Elevated plasmacytokine levels have been related to the development of systemic inflammatory response syndrome, sepsis, and organ dysfunction.

TNF- α has been associated with patients at high-risk for sepsis after abdominal surgery and is a predictor of mortality [8, 9]. Qui P et al. [10] investigated the therapeutic role of anti-TNFagents in a meta-analysis study. They searched 15 studies and concluded that anti-TNF agents produced a modest but significant decrease in




FIGURE 2. Lung tissue of Group I (A), Group II (B), and Group III (C). Leucocyte infiltration can be seen in Group I. A decrease in leucocyte infiltration can be seen in Groups II and III.

TABLE 3. Comparison of TNF, IL-1beta, IL-6 and CRP levels parameters according to the groups

	Group	Mean	Median	St. Dev.	Min.	Max.	p	p1	p2	p3
TNF	I	248.8	251.1	21.4	213.9	275.4	<0.05*	0.073	0.003	0.038
	II	225.1	226.8	13.0	209.8	237.0				
	III	200.4	196.3	16.9	185.1	229.9				
IL-1beta	I	1936.2	1933	216.8	1631.2	2271.0	0.041*	0.214	0.419	0.914
	II	1866.9	1859.2	116.7	1758.8	1990.7				
	III	1657.3	1677.3	55.2	1556.3	1710.9				
IL-6	I	9.3	9.4	1.0	7.9	10.6	0.424*	-	-	-
	II	10.6	10.3	1.7	9	12.6				
	III	9.9	9.9	1.3	8.0	12.0				
CRP	I	200.7	202.7	11.1	175.6	210.9	0.011*	0.808	0.001	0.114
	II	196.3	206.5	43.9	134.9	327.4				
	III	154.1	163.4	19.3	128.0	170.5				

*Kruskal Wallis p value; 1I vs. II, 2I vs. III, 3I vs. II (Mann-Whitney U p value).

mortality in patients with sepsis.

High levels of CRP are caused by infections and many inflammatory diseases. The relationship between intra-abdominal infections and CRP is well established [11]. K. Mulari analyzed 66 patients with secondary peritonitis due to gastrointestinal tract perforation and investigated risk factors for hospital mortality. It was concluded that elevated CRP levels and high Mannheim peritonitis index score in the early postoperative phase had a prognostic significance [12].

Peritoneal lavage is described as the washing of the peritoneal space with high volumes of saline. Different agents have been used for peritoneal lavage in the literature. Povidone–iodine (PVI) and saline are well-known agents that are used for this purpose. Araujo ID et al. investigated peritoneal lavage with PVI in rats with exper-

imental peritonitis. Lavage of the peritoneal cavity with PVI demonstrated no beneficial effect in local control of peritonitis [13]. Camargo M et al, showed a beneficial effect of peritoneal lavage with bupivacaine in rats with fecal peritonitis [14]. Peritoneal irrigation was performed in cases of severe pancreatitis and morbidity and mortality rates decreased in this patient group [15]. Coumarin, which has an immunostimulant effect, was used in peritonitis without any beneficial effect [16, 17]. Peritoneal irrigation performed with a local anesthetic agent, 0.2% ropivacaine, in rats with fecal peritonitis decreased the histologic changes that occurred due to inflammation [18].

In our study, Groups II and III underwent the proposed therapeutic process (peritoneal lavage with isotonic solution and mesalazine). The effects of peritoneal lavage with isotonic solution and mesalazine was exam-

ined objectively. The mean CRP level was found statistically significantly lower in Group III than in the other groups. TNF α and IL-1 β levels were also lower in Group III. The decrease in TNF- α levels in Group III was statistically significant. We examined the lung tissues to show the systemic effect of intra-abdominal infection. The degree of inflammatory reaction was found lower in Group III, indicating that lavage using mesalazine may have a protective effect on the lung tissue.

CONCLUSION

In rats with intra-abdominal sepsis, TNF- α and CRP levels significantly decreased in the mesalazine group. There were also decrease in leukocyte infiltration in the lungs of rats treated with mesalazine. So, it may be suggested that the use of mesalazine in clinical practice may decrease the sepsis-related morbidity and mortality.

Conflict of Interest: The authors declare no conflict of interest.

Financial Disclosure: The authors declared that this study has received no financial support.

Authorship Contributions: Concept – A.S.; Design – A.S.; Supervision – B.K.; Materials – A.S., M.S.T., L.T.S., B.K.; Data collection &/or processing – L.T.S.; Analysis and/or interpretation – M.S.T., L.T.S., A.S.; Writing – A.S., M.S.T., L.T.S., A.S., B.K.; Critical review – A.S.

REFERENCES

1. Angus DC, Linde-Zwirble WT, Lidicker J, Clermont G, Carcillo J, Pinsky MR. Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated costs of care. *Crit Care Med* 2001;29:1303–10. [\[CrossRef\]](#)
2. Palmiere C, Augsburg M. Markers for sepsis diagnosis in the forensic setting: state of the art. *Croat Med J* 2014;55:103–14. [\[CrossRef\]](#)
3. Hatamkhani S, Karimzadeh I, Elyasi S, Farsaie S, Khalili H. Carnitine and sepsis: a review of an old clinical dilemma. *J Pharm Pharm Sci* 2013;16:414–23. [\[CrossRef\]](#)
4. Campieri M, Lanfranchi GA, Bazzocchi G, Brignola C, Sarti F, Franzin G, et al. Treatment of ulcerative colitis with high-dose 5-aminosalicylic acid enemas. *Lancet* 1981;2:270–1. [\[CrossRef\]](#)
5. Svartz M. The treatment of 124 cases of ulcerative colitis with salazopyrine and attempts of desensibilization in cases of hypersensitiveness to sulfa. *Acta Med Scand* 1948;131:465–72. [\[CrossRef\]](#)
6. Koelink PJ, Hawinkels LJ, Wiercinska E, Sier CF, ten Dijke P, Lamers CB, et al. 5-Aminosalicylic acid inhibits TGF- β 1 signalling in colorectal cancer cells. *Cancer Lett* 2010;287:82–90. [\[CrossRef\]](#)
7. Sands KE, Bates DW, Lanken PN, Graman PS, Hibberd PL, Kahn KL, et al. Academic Medical Center Consortium Sepsis Project Working Group. Epidemiology of sepsis syndrome in 8 academic medical centers. *JAMA* 1997;278:23–40. [\[CrossRef\]](#)
8. Davis MG, Hagen PO. Systemic inflammatory response syndrome. *Br J Surg* 1997;84:920–35. [\[CrossRef\]](#)
9. Oda S, Hirasawa H, Shiga H, Nakanishi K, Matsuda K, Nakamura M. Sequential measurement of IL-6 blood levels in patients with systemic inflammatory response syndrome (SIRS)/sepsis. *Cytokine* 2005;29:169–75. [\[CrossRef\]](#)
10. Qiu P, Cui X, Sun J, Welsh J, Natanson C, Eichacker PQ. Antitumor necrosis factor therapy is associated with improved survival in clinical sepsis trials: a meta-analysis. *Crit Care Med* 2013;41:2419–29.
11. Kørner H, Nielsen HJ, Søreide JA, Nedrebø BS, Søreide K, Knapp JC. Diagnostic accuracy of C-reactive protein for intraabdominal infections after colorectal resections. *J Gastrointest Surg* 2009;13:1599–606.
12. Mulari K, Leppäniemi A. Severe secondary peritonitis following gastrointestinal tract perforation. *Scand J Surg* 2004;93:204–8. [\[CrossRef\]](#)
13. Araujo ID, Grossi GC, Diniz SO, Nunes TA, Braga EA, Cardoso VN. Effects of the povidone-iodine (PVPI) in treatment of bacterial peritonitis induced in rats. *Acta Cir Bras* 2010;25:322–7. [\[CrossRef\]](#)
14. Camargo MG, Fagundes JJ, Leal RE, Ayrizono Mde L, Rossi DH, Oliveira Pde S, et al. Influence of the peritoneal lavage with bupivacaine on the survival and resistance of colonic anastomoses performed under fecal peritonitis in rats. *Acta Cir Bras* 2013;28:783–7. [\[CrossRef\]](#)
15. Matsumoto K, Miyake Y, Nakatsu M, Toyokawa T, Ando M, Hirohata M, et al. Usefulness of early-phase peritoneal lavage for treating severe acute pancreatitis. *Intern Med* 2014;53:1–6. [\[CrossRef\]](#)
16. Sortini D, Feo CV, Maravegias K, Carcoforo P, Pozza E, Liboni A, et al. Role of peritoneal lavage in adhesion formation and survival rate in rats: an experimental study. *J Invest Surg* 2006;19:291–7. [\[CrossRef\]](#)
17. Creagh TA, Leahy AL, McNamara E, Bouchier-Hayes DJ. Local beneficial effect of coumarin in experimental peritonitis. *Ir J Med Sci* 1991;160:385–6. [\[CrossRef\]](#)
18. Brocco MC, Gomez RS, Paulo DN, Almeida CE, Baptista JF. Histological features of peritoneal lavage with ropivacaine in rats with fecal peritonitis. *Acta Cir Bras* 2012;27:193–9. [\[CrossRef\]](#)