

Bullous drug eruption with low dose methotrexate

 Tuba Yuce Inel

Division of Rheumatology, Department of Internal Diseases, Dokuz Eylul University Faculty of Medicine, Izmir, Turkey

Cite this article as: Yuce Inel T. Bullous drug eruption with low dose methotrexate. *North Clin Istamb* 2021;8(6):617–618.

A 71-year-old woman was referred by an ophthalmologist who treated her corneal melt. Although she had erosive rheumatoid arthritis, she was not followed up for 5 years. After the diagnosis was confirmed by clinical and radiological, methotrexate (MTX) 7.5 mg/week was started because of the arthritis in her right wrist and metacarpophalangeal joints. Two days later, bullous lesions developed on the dorsal aspect of the left foot, and behind the left leg (Fig. 1). She did not have a fever or mucosal lesions. No drug-related cytopenia was observed. Levels of albumin, creatinine, and liver function tests were normal, proteinuria was not detected. Anti-nuclear antibody, anti-neutrophil cytoplasmic antibody, complements (C3–C4), and hepatitis serology were negative. MTX treatment was discontinued. Skin biopsy was performed for the differential diagnosis. Direct immunofluorescence examination of skin biopsy was negative and reported as bullous drug eruption. After the intact bullae were opened with a sterile injector, they were dressed with fusidic acid and ethacridine lactate. Meropenem and linezolid antibiotherapy were given to the patient who developed wound infection. Two weeks later, healthy re-epithelization was observed at the wound site.

Cutaneous side effects of MTX are secondary to hypersensitivity reactions or induction of apoptotic cell death. Bullous acral erythema may develop with high dose MTX treatment which occurs 1–21 days after chemotherapy administration [1]. This reaction is usually seen at high doses, but also there are few reports indicating that it is also seen at low doses [2]. It should be kept in mind that bullous drug eruption may be seen with low dose MTX, possibly as a result of an idiosyncratic reaction.




FIGURE 1. Bullous lesions on the left leg and dorsal aspect of the foot.

Informed Consent: Written informed consent was obtained from the patient for the publication of the case report and the accompanying images.

Received: September 21, 2020 *Accepted:* November 04, 2020 *Online:* December 31, 2021



Correspondence: Tuba YUCE INEL, MD. Dokuz Eylul Universitesi Tip Fakultesi, Ic Hastaliklari Anabilim Dalı, Romatoloji Bilim Dalı, Izmir, Turkey.

Tel: +90 232 412 37 27 e-mail: dr.tubayuce@yahoo.com

© Copyright 2021 by Istanbul Provincial Directorate of Health - Available online at www.northclinist.com

REFERENCES

1. Kuruvila S, Dalal M, Sivanesan B. Bullous variant of acral erythema due to methotrexate. *Indian J Dermatol Venereol Leprol* 2006;72:440–2. [\[CrossRef\]](#)
2. Sako EY, Famenini S, Wu JJ. Bullous drug eruption in a patient with psoriasis after a test dose of methotrexate. *J Am Acad Dermatol* 2013;69:e264–5. [\[CrossRef\]](#)