

Keywords: Extranodal natural killer/T-cell lymphoma, Non-Hodgkin lymphoma

Anahtar Sözcükler: Ekstranodal natural killer/T-hücreli lenfoma, Hodgkin dışı lenfoma

Authorship Contributions

Concept: Esra Sarıbacak Can, Design: Esra Sarıbacak Can, Data Collection or Processing: Esra Sarıbacak Can, Analysis or Interpretation: Esra Sarıbacak Can, Harika Okutan, Literature Search: Esra Sarıbacak Can, Murat Albayrak, Writing: Esra Sarıbacak Can, Ünsal Han.

Conflict of Interest: The authors of this paper have no conflicts of interest, including specific financial interests, relationships, and/or affiliations relevant to the subject matter or materials included.

References

1. Roschewski M, Wilson WH. EBV-associated lymphomas in adults. *Best Pract Res Clin Haematol* 2012;25:75-89.
2. Niedobitek G. Epstein-Barr virus infection in the pathogenesis of nasopharyngeal carcinoma. *Mol Pathol* 2000;53:248-254.
3. Al-Hakeem DA, Fedele S, Carlos R, Porter S. Extranodal NK/T-cell lymphoma, nasal type. *Oral Oncol* 2007;43:4-14.
4. Jaccard A, Hermine O. Extranodal natural killer/T-cell lymphoma: advances in the management. *Curr Opin Oncol* 2011;23:429-435.
5. Liu QF, Wang WH, Wang SL, Liu YP, Huang WT, Lu N, Zhou LQ, Ouyang H, Jin J, Li YX. Immunophenotypic and clinical differences between the nasal and extranasal subtypes of upper aerodigestive tract natural killer/T-cell lymphoma. *Int J Radiat Oncol Biol Phys* 2014;88:806-813.
6. Liang R. Diagnosis and management of primary nasal lymphoma of T-cell or NK-cell origin. *Clin Lymphoma* 2000;1:33-38.
7. Hasserjian RP, Harris NL. NK-cell lymphomas and leukemias: a spectrum of tumors with variable manifestations and immunophenotype. *Am J Clin Pathol* 2007;127:860-868.
8. Kwong YL. Natural killer-cell malignancies: diagnosis and treatment. *Leukemia* 2005;19:2186-2194.



Address for Correspondence/Yazışma Adresi: Esra SARIBACAK CAN, M.D.,
Dışkapı Yıldırım Beyazıt Research and Training Hospital,
Clinic of Hematology, Ankara, Turkey
E-mail : esracanercan@hotmail.com

Received/Geliş tarihi: April 27, 2015
Accepted/Kabul tarihi: September 02, 2015

DOI: 10.4274/tjh.2015.0169

T-Cell Lymphoma Presenting with Auricular and Parotid Gland Involvement

Aurikula ve Parotis Bezi Tutulumu ile Seyreden T-Hücreli Lenfoma

Birgül Öneç¹, Alper Koç², Elif Nisa Ünlü³, İlhan Ünlü⁴, Hüseyin Yaman⁴, Durdu Mehmet Köş²

¹Düzce University Faculty of Medicine, Department of Hematology, Düzce, Turkey

²Düzce University Faculty of Medicine, Department of Internal Medicine, Düzce, Turkey

³Düzce University Faculty of Medicine, Department of Radiology, Düzce, Turkey

⁴Düzce University Faculty of Medicine, Department of Otorhinolaryngology, Düzce, Turkey

To the Editor,

The external auditory canal is an unusual presenting site for lymphomas, with only a few case reports in the literature [1,2]. Malignant lymphomas arising from the salivary glands are also uncommon, accounting for approximately 5% of extranodal lymphomas, and the majority of them are of B-cell lineage. Primary salivary gland T-cell lymphomas are extremely rare [3,4,5].

A 63-year-old man was admitted with swelling of the left side of his face and left auricle. Considered as an infection, it was empirically treated with systemic and topical antibiotics. Examination by an otorhinolaryngologist revealed a suppurative

lesion that consisted of ulcerated areas and granulation tissue in the external auditory canal and preauricular region in addition to edema (Figure 1A). A lobulated mass lesion of 47x39 mm arising from the left parotid and extending to the left auditory canal was detected in computed tomography (CT) with accompanying lymph nodes in the left subauricular and cervical localization. Fine-needle biopsy was nondiagnostic and incisional biopsy found only severely active chronic inflammation. Finally, deep excisional biopsy revealed CD3-, CD5-, and CD30-positive and S100-negative lymphoid cells diffusely infiltrating the dermis. The case was considered as stage 2 peripheral T-cell lymphoma not otherwise specified peripheral T-cell lymphoma-not otherwise specified (PTCL-NOS) and CHOP (cyclophosphamide,



Figure 1. A) A suppurative lesion that consisted of ulcerated areas and granulation tissue was observed in the external auditory canal and preauricular region. B) Significant regression of lesion after 4 cycles of CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) treatment.

doxorubicin, vincristine, prednisone) protocol was started. Despite clinical improvement in the lesions (Figure 1B), CT revealed progression after the fourth cycle. The second-line treatment is ongoing with DHAP (dexamethasone, high-dose ara-C, cisplatin) at the 8th month of follow-up.

The auricula, external auditory canal, and parotid glands are unusual locations for T-cell lymphoma. Presentations of lymphomas are indistinguishable from other swellings of the auricle or parotid gland and therefore a high index of suspicion should be maintained in patients who present with presumptive cutaneous infections that do not respond to antibiotic therapy in these locations. A suppurative auricular lesion suggests an infectious disease rather than a lymphoma, but primary cutaneous lymphomas and cutaneous manifestations of lymphomas must be kept in mind. Early excisional biopsy may prevent excessive waste of time with unnecessary antibiotherapies. Parotid lymphomas are most likely to be B-cell non-Hodgkin lymphoma, but non-B-cell lymphomas have a more aggressive course in all salivary gland lymphomas. Both B-cell and T-cell lymphomas share many morphological similarities; therefore, immunohistochemical analysis is required

for proper assignment of lineage of salivary gland lymphomas [5]. Reactive lymphoid infiltrate located in the periphery of the lymphoma may contribute to the delay of diagnosis and larger biopsy samples are needed.

Keywords: Parotid gland, T-Cell lymphoma, Auricula

Anahtar Sözcükler: Parotis bezi, T-Hücreli lenfoma, Aurikula

Authorship Contributions

Concept: Birgül Öneç, Design: Birgül Öneç, Alper Koç, Data Collection or Processing: Birgül Öneç, Alper Koç, Analysis or Interpretation: Birgül Öneç, Durdu Mehmet Köş, Elif Nisa Ünlü, Literature Search: Birgül Öneç, Hüseyin Yaman, İlhan Ünlü, Writing: Birgül Öneç, Alper Koç.

Conflict of Interest: The authors of this paper have no conflicts of interest, including specific financial interests, relationships, and/or affiliations relevant to the subject matter or materials included.

References

1. Marçal N, Campelos S, Dias L, Gonçalves M, Pereira G, Godinho T. Primary cutaneous CD30-positive anaplastic large-cell lymphoma of the external auditory canal. *Ear Nose Throat J* 2012;91:10-12.
2. González Delgado A, Argudo Marco F, Sánchez Martínez N, Sprekelsen Gassó C. T cell non-Hodgkin's lymphoma of the external auditory canal. *Acta Otorrinolaringol Esp* 2008;59:200-201.
3. Yanagawa N, Osakabe M, Furuse H, Maeda K, Tamura G. Peripheral T-cell lymphoma arising from an intraglandular lymph node in the parotid gland: a case report and literature review. *Pathol Int* 2012;62:60-64.
4. Metikurke SH, Krishnappa R, Ramachar SM, Arora I. Primary malignant lymphoma of the parotid gland. *J Cancer Res Ther* 2012;8:641-643.
5. Chan JK, Tsang WY, Hui PK, Ng CS, Sin VC, Khan SM, Siu LL. T- and T/ natural killer-cell lymphomas of the salivary gland: a clinicopathologic, immunohistochemical and molecular study of six cases. *Hum Pathol* 1997;28:238-245.



Address for Correspondence/Yazışma Adresi: Alper KOÇ, M.D.,
Düzce University Faculty of Medicine, Department of Internal Medicine,
Düzce, Turkey
E-mail : alperkoc44@hotmail.com

Received/Geliş tarihi: May 25, 2015
Accepted/Kabul tarihi: August 17, 2015

DOI: 10.4274/tjh.2015.0217