

Heart Failure 2019 Insights from the National Society of Cardiology journals

**Plamen Gatzov M.D.¹, Jean-Jacques Monsuez M.D.², Gergely Agoston M.D.³, Michael Aschermann M.D.⁴,
Hala Mahfouz Badran M.D.⁵, Ariel Cohen M.D.⁶, Kurt Huber M.D.⁷, Evgeny Shlyakhto M.D.⁸,
Dilek Ural M.D.⁹, Ignacio Ferreira-Gonzalez M.D.¹⁰, Fernando Alfonso M.D.¹¹**

¹Editor-in-Chief of Bulgarian Cardiology Journal, Bulgaria; Spain

²Editor-in-Chief of Archives des Maladies du Coeur et des Vaisseaux Pratique, France

³Editor-in-Chief of Cardiologia Hungarica, Hungary

⁴Editor-in-Chief of Cor et Vasa, Czech Republic

⁵Editor-in-Chief of Egyptian Heart Journal, Egypt

⁶Editor-in-Chief of Archives of Cardiovascular Diseases, France

⁷Editor-in-Chief of Austrian J Kardiologie, Austria

⁸Editor-in-Chief of Russian Journal of Cardiology, Russia

⁹Editor-in-Chief of Archives of The Turkish Society of Cardiology, Turkey

¹⁰Editor-in-Chief of Revista Española de Cardiología, Spain

¹¹Chairman of The Editors Network of the European Society of Cardiology, Spain

Most studies on heart failure (HF) management published in 2019 by high-ranking impact factor international journals focus on drug therapy. This included administration of sacubitril-valsartan with initiation during the index admission and the benefits of SGLT2 inhibitors in reducing cardiovascular mortality and HF. Most of these studies, targeting a broad readership, fail to characterize important local issues.

Improvement in HF management needs to take into account specificities from different European Society of Cardiology (ESC) member countries. This approach may be achieved and disseminated to cardiologists by the National Society of Cardiology Journals (NSCJ). During the ESC Congress 2019, the ESC Editors' Network started an initiative intended

to boost dissemination of cardiology research published in the NSCJ by summarizing in a review paper the evidence gathered in selected areas. The ESC Editors Network members decided the first topic of such a review to be publications in the field of HF.

Epidemiology

Inequalities in the prevalence of risk factors, cardiovascular disease burden, cardiovascular mortality, and implementation of some therapeutic methods (coronary interventions, device implantations, and cardiac surgery) among the ESC member countries have recently been shown in the ATLAS study. These shortcomings depend on socioeconomic factors and affect more predominantly middle-income than high-income countries. In a single-centre observa-

Reproduced from: Gatzov P, Monsuez JJ, Agoston G, Aschermann M, Badran HM, Cohen A, et al. Heart Failure 2019 Insights from the National Society of Cardiology journals. European Heart Journal 2021;42(6):557-9. doi:10.1093/eurheartj/ehaa918, by permission of Oxford University Press on behalf of the European Society of Cardiology.

© The Author(s) 2020.

All rights reserved; no part of this publication may be produced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without prior written permission of the Publishers.

For Permissions, please email: journals.permission@oup.com

The opinions expressed in the Journal item reproduced as this reprint are those of the authors and contributors, and do not necessarily reflect those of the European Society of Cardiology, the editors, the editorial board, Oxford University Press or the organization to which the authors are affiliated.

Oxford University Press, OPL, and the European Society of Cardiology are not responsible or in any way liable for the accuracy of the reprint, for any errors, omissions, or inaccuracies, or for any consequences arising therefrom. The Turkish Society of Cardiology is solely responsible for this reprint.

Correspondence: Prof. Plamen Gatzov, MD, PhD, DSc, FESC,
Medical University of Plevan, 1, St. Kl. Ohridski Street, 5800 Plevan, Bulgaria.
e-mail: plamengatzov@yahoo.com



tional study of 1006 patients admitted to a coronary care unit in Egypt, Badran et al.^[11] estimated the prevalence of HF by gender and preserved or reduced left ventricular ejection fraction (LVEF). They reported a higher prevalence of HF and a higher incidence of HF with reduced ejection fraction (HFrEF) among women. Female patients were older, more likely to be obese, with more co-morbidities, had less acute coronary syndromes and required fewer coronary interventions, but had a prognosis similar to men.

Nationwide information on mortality and readmissions in HF patients is also of interest. A retrospective analysis from Spain aimed to identify factors associated with in-hospital mortality and readmissions in 77 652 HF patients.^[12] In-hospital mortality was 9.2%, rising to 14.5% at 1 year. The 1-year cardiovascular readmission rate was 33%. Risk factors associated with mortality were stroke, metastatic cancer, cardiorespiratory failure, shock, and acute myocardial infarction. Risk-standardized mortality rates were lower among patients discharged from high-volume hospitals. Importantly, the availability of a cardiology department at the hospital was associated with better outcomes.

Specific causes of heart failure

Calcific aortic valve disease (CAVD) is a common disorder which may progress while remaining clinically unrecognized. However, the progression mechanisms remain unknown. In a single-centre study from Bulgaria, Tomova et al.^[13] tested the hypothesis that the polymorphism rs10455872 at the lipoprotein (a) (Lp(a)) gene locus increases the risk of CAVD. In a comparison of 108 CAVD patients with 38 controls, they reported that the presence of ≥ 1 mutant allele of the gene was associated with a four-fold greater risk for CAVD. A report from the Austrian Journal of Cardiology by Kauffmann et al.^[14] demonstrated that the diagnosis of transthyretin-amyloidosis has recently improved significantly with structural screening by magnetic resonance imaging (MRI). The coexistence of CAVD and transthyretin-amyloidosis has substantial implications. The study suggested the value of the electrocardiogram, echocardiography, MRI, technetium radionuclide imaging, and endomyocardial biopsy in these patients.^[14] Subclinical myocardial involvement is common in systemic sclerosis (SSc) and is associated with HF and a worse prognosis. In a study of 73 SSc patients from Hungary, Vertes et al.^[15]

tested the 2D-speckle-tracking-derived global longitudinal strain (GLS) for the detection of early myocardial involvement. Lower GLS values were found in patients compared with gender- and age-matched healthy controls. GLS values correlated with the duration of the disease from the onset of Raynaud's phenomenon, from the first non-Raynaud symptoms, and with the New York Heart Association (NYHA) functional class.

Pathogenesis

In a prospective Russian study including 297 patients, Lelyavina et al.^[16] reported the potential for differentiation, regeneration, and growth of satellite skeletal muscle precursor cells obtained from patients with HFrEF. The studied parameters were similar to those found in healthy donors. This may explain why walking >1.5 h/day induces more physiological reverse myocardial remodelling than aerobic training.

Diagnosis

In another Russian study, Vdovenko et al.^[17] compared 80 patients (NYHA Class I-IIa) with chronic HF-pEF with 30 healthy controls by using the 6-minute walk-test and echocardiography. All patients had diastolic dysfunction (60 abnormal relaxation patterns and 20 pseudo normal patterns) with reduced global and segmental LV strain. The impact of HF on other body organs has been analysed by Içen et al.^[18] from Turkey assessing liver stiffness (LS) in HF patients. Liver stiffness estimated using an ElastPQ technique was increased in patients with more advanced NYHA class. A higher LS was associated with higher right ventricular myocardial performance index, regurgitation pressure gradient, NT-proBNP, and aspartate aminotransferase levels. Assuming that the SYNTAX score is not just a measure of the severity but also the complexity of coronary artery disease Öztürk et al.^[19] analysed the degree of coronary atherosclerosis, estimated by SYNTAX score and myocardial viability in Turkish patients with ischaemic cardiomyopathy. Patients without viability had a significantly higher SYNTAX score compared with those with viable myocardium.

Treatment

Clinical implications of HF associated with valvular heart disease has been reported by several NSCJ. Transcatheter aortic valve implantation (TAVI) is an alternative to surgical aortic replacement for symp-

omatic severe aortic stenosis. Indications are now rapidly expanding towards patients at lower surgical risk. Generalization of the transfemoral vascular approach, technological advances, and increased operator skills have resulted in higher rates of procedural success and improved long-term survival. However, data on the incidence of readmission for HF after successful TAVI are scarce.

A French study on 1139 patients, by Guedeney et al.^[10] reported that readmission for HF occurs in 1/10 patients after TAVI and suggests a strong risk factor for mortality. The main risk factors for HF readmission were LVEF $\leq 35\%$, chronic pulmonary disease, chronic kidney disease, diabetes, and atrial fibrillation. Along with international multicentre trials on the percutaneous mitral repair of functional mitral regurgitation associated with HF, national registries provide valuable real-life results in unselected patients which may inform clinical decisionmaking at a local level.

Benak et al. reported a cohort of 30 MitraClip implantations in Czech patients with dilated cardiomyopathy and severe functional mitral regurgitation. Procedural success was 97% with no 90-day mortality. At 1 year, significant improvements in functional class, and quality of life scores were reported, associated with a reduction of LV myocardial mass, an increase in systolic and diastolic arterial pressure and a mortality rate of 10%.

Conclusion

Instead of publications in high-ranking impact factor international journals, NSCJ cover a wide spectrum of diagnostic and therapeutic modalities taking into account the national specifics of HF. In contrast, studies published in NSCJ are often single-centre and observational. However, information on HF strategies at a national level are useful in implementing ESC clinical practice guidelines and in optimizing HF patients' care.

Conflict of interest: None.

REFERENCES

1. Badran HM, Elgharably MA, Faheem N. Clinical characteristics and in-hospital outcome of heart failure in women: a single center registry from Egyptian cardiac care unit. *Egypt Heart J* 2019;71:167. [\[Crossref\]](#)
2. Martínez Santos P, Bover Freire R, Esteban Fernández A, Bernal Sobrino JL, Fernández Pérez C, Elola Somoza FJ, et al. In-hospital mortality and readmissions for heart failure in Spain. A Study of Index Episodes and 30-Day and 1-year Cardiac Readmissions. *Rev Esp Cardiol (Engl Ed)* 2019;72:998-1004. [\[Crossref\]](#)
3. Tomova V, Alexandrovna M, Atanasova M, Rashev T, ceko-va M. The polymorphism RS10455872 at the lipoprotein (a) gene locus enhances the risk of aortic valve disease. *Bulg Cardiol J* 2019;4:45-50. [\[Crossref\]](#)
4. Kaufmann CC, Hennenberg J, Bonderman D, Kriz R, Koch J, Huber K, Jager B. Transthyretin amyloidosis - update 2019. *Austr J Kardiol* 2019;26:112-6.
5. Vertes V, Nogradi A, Porpaczy A, Minier T, Czirjak L, Komocsi A, et al. Left ventricular global longitudinal strain is impaired in systemic sclerosis and shows a significant correlation with the functional capacity of the patients. *Cardiol Hung* 2019;49:12-16.
6. Lelyavina TA, Sitnikova MY, Galenko VL, Kozlov PS, Bortsova MA, Demchenko EA, et al. The role of muscle tissue in the pathogenesis of chronic heart failure-the potential of exposure (FORMA study). *Russ J Cardiol* 2019;2:58-65. [\[Crossref\]](#)
7. Vdovenko DV, Libis RA. Assessment of the functional state of left heart in patients with chronic heart failure with preserved ejection fraction. *Russ J Cardiol* 2019;2:26-30. [\[Crossref\]](#)
8. Içen YK, Demirtas AO, Koç AS, Sumbul HE, Koc M. Liver stiffness value obtained with ElastPQ ultrasound increases with NYHA class in chronic heart failure patients and reduced ejection fraction. *Turk Kardiyol Dern Ars* 2019;47:281-3. [\[Crossref\]](#)
9. Öztürk S, Gürbuz AS, Kirma C. Relationship between SYNTAX score and myocardial viability in ischemic cardiomyopathy. *Turk Kardiyol Dern Ars* 2019;47:350-6. [\[Crossref\]](#)
10. Guedeney P, Huchet F, Manigold T, Rouanet S, Balagny P, Leprince P, et al. Incidence of risk factors for and impact of readmission for heart failure after successful transcatheter aortic valve implantation. *Arch Cardiovasc Dis* 2019;112:765-72. [\[Crossref\]](#)