

Management of tracheal stenosis after corrosive acid ingestion: A case report

Hidir Esme, M.D., Hasan Doğan, M.D.

Department of Thoracic Surgery, University of Health Sciences, Konya Training and Research Hospital, Konya-Turkey

ABSTRACT

Although corrosive injury of the digestive tract is a well-known clinical entity, damages of the airway and a critically life-threatening condition, have not been clearly documented. Tracheal stenosis is very rare associated with corrosive acid ingestion. We report the case of a 4-year-old girl child who presented to the emergency department three weeks after accidentally drinking an acidic cleaning agent stored in an unlabeled bottle. Rigid bronchoscopy was administered to observe the stenosis. She was treated by serial dilation, repair of tracheal laceration, and placement of a temporary polyurethane-coated nitinol stent. Careful and accurate stent placement may provide significant and life-saving airway improvement as observed in the presenting pediatric case.

Keywords: Corrosive injury; management; tracheal stenosis.

INTRODUCTION

Corrosive acid poisoning commonly results in chemical injuries to the upper gastrointestinal tract. Corrosive mucosal erosion of the larynx and trachea may occur if the patient aspirates acid. Although corrosive injury of the digestive tract is a well-known clinical entity, such as damages of the airway, a critically life-threatening condition, has not been clearly documented.^[1] Tracheal stenosis is very rare and associated with corrosive acid ingestion. Caustic ingestion results in thrombosis of small vessels with inflammation, the formation of granulation tissue with subsequent collagen deposition and fibrosis, thus stricture formation.^[2]

In this case study, we report an incident of upper respiratory system corrosion after aspiration of the caustic acid agent. The patient survived the severe burns of the tracheal tract and the tracheal stenosis by dilation procedures and surgical repair of the iatrogenic tracheal laceration and treatment of the tracheal stenosis with a temporary polyurethane-coated nitinol stent. These impressive clinical features are presented and the management of the caustic respiratory injury is discussed.

CASE REPORT

A 4-year-old girl child presented to the emergency department three weeks after accidentally drinking an acidic cleaning agent stored in an unlabeled bottle. On the day of the incident, the patient presented to an outside hospital where she was admitted for an upper endoscopy of the esophagus as well as laryngoscopy, which were both negative for acute injury. An initial chest X-ray taken the day of the incident was also normal. After two days of observation, the patient was discharged.

When patient was admitted to our department, she continued to have a sore throat and upper chest pain with associated shortness of breath with inspiratory stridor. The patient denied fever, chills, weakness, bloody stools, or upper and lower gastrointestinal upset. On presentation, the patient vital signs were as follows: blood pressure of 98/57 mmHg, the pulse of 121, respiratory rate of 29, the temperature of 36.7°C and oxygen saturation of 90% on room air. Her physical exam was significant for pharyngeal and uvula erythema and edema without ulcers. Her lung exam demonstrated slightly dimin-

Cite this article as: Esme H, Doğan H. Management of tracheal stenosis after corrosive acid ingestion: A case report. *Ulus Travma Acil Cerrahi Derg* 2021;27:377-379.

Address for correspondence: Hidir Esme, M.D.

Sağlık Bilimleri Üniversitesi, Konya Eğitim ve Araştırma Hastanesi, Göğüs Cerrahi Kliniği, Konya, Turkey

Tel: +90 332 - 322 28 19 E-mail: drhesme@hotmail.com

Ulus Travma Acil Cerrahi Derg 2021;27(3):377-379 DOI: 10.14744/tjtes.2020.21805 Submitted: 31.01.2019 Accepted: 10.03.2020

Copyright 2021 Turkish Association of Trauma and Emergency Surgery



ished breathe sounds bilaterally with noted increased work of breathing. His abdomen, including the epigastric region, was soft without rebound or guarding. The remainder of his exam was normal.

An anterior chest X-ray was taken with showed clear lungs fields without effusion, infiltrate or aspiration, normal mediastinum, and no lymphadenopathy. The computed tomographic scan was significant for tracheal stenosis (Fig. 1a). Rigid bronchoscopy was administered to observe the stenosis. After inspection and measurement, serial dilation was done with successively larger bronchoscopes. The patient recovered well after bronchoscopy and she was discharged. After 20 days, tracheal stenosis was repeated. At that time, rigid bronchoscopy revealed more than 70% luminal narrowing. Dilatation with rigid bronchoscopy was performed. Postoperative bilateral pneumothorax, subcutaneous emphysema developed (Fig. 1b). This time, bronchoscopy revealed partial tracheal laceration in the thoracic part of the trachea. The tracheal injury consisted of a 2 cm tear extending from approximately 3 cm distal to the vocal cords and ending 1 cm proximal to the carina. Tracheal perforation was repaired with right thoracotomy. Bilateral chest drains were inserted and kept for seven days. The inspiratory stridor continued in the patient. A 10-mm x 4-cm nitinol fully covered stent was placed in the trachea to just above the carina (Fig. 1c). The patient was discharged ten days later with good chest expansion and had been doing well for more than two months without any subsequent problems. Consent was obtained from the patient's family for this study.

DISCUSSION

The accidental ingestion of easily available corrosive substances is a significant social problem. Although it rarely causes mortality, its morbidity lasts a lifetime. Corrosive injury of the upper gastrointestinal tract is a common clinical entity. Airway aspiration, instead of ingestion of the caustic substance, as in the presenting case, is another mechanism

of pulmonary injury, which causes direct burning of the respiratory system.^[1] Airway bleeding and obstruction with tissue slough are the early clinical clue of caustic aspiration. A key to alkaline injury is that it causes liquefactive tissue necrosis leading to the dissolution of cellular components and saponification of fatty tissues resulting in a liquid-gel amalgamation of dissolved cells and connective tissue. Ingestion of acidic media results in immediate denaturing of proteins which limits proteolysis of cellular constituents and leads to localized eschar formation, which limits further tissue damage.^[3] Our patient presented to our department three weeks after accidentally drinking an acidic cleaning agent. We have linked the late manifestation of respiratory symptoms to the ingestion of corrosive acid agents.

Tracheal stenosis is uncommon in the pediatric age group. It may be due to congenital atresia, tracheomalacia or acquired stenosis. Acquired stenosis is mostly due to prolonged endotracheal intubation, faulty tracheostomy or external trauma.^[4] Tracheal stenosis is very rare associated with corrosive ingestion. Tracheal stenosis produces symptoms of dyspnea, stridor, and obstructive pneumonia and is frequently life-threatening, with patients having impending suffocation. Bronchoscopic management is the first step in providing a diagnosis, stabilizing the obstructed airway, and evaluating resectability. Tracheal stenosis is dilated with esophageal bougies, the bronchoscope, or appropriately sized angioplasty balloons. After inspection and measurement, serial dilation is done with successively larger bronchoscopes. Steady rotating pressure with the blunt-tipped Jackson bronchoscopes is provided passage with minimal mucosal trauma and risk of perforation. The procedure is repeated with the next larger bronchoscope until an adequate airway caliber had been established. If the lesion is too stenotic to accept the 3.5-mm Jackson bronchoscope, then esophageal bougies are used to enlarging the airway enough to allow bronchoscopic dilation. Pneumatic or hydrostatic balloon dilation with angioplasty balloons are used whenever it is necessary to dilate larger than the 8 to 9 mm possible with bronchoscopic dilation.^[5]




Figure 1. (a) The computed tomographic scan showing tracheal stenosis as the result of acid ingestion. (b) Anterior-posterior chest X-ray demonstrating bilateral pneumothorax, subcutaneous emphysema after bronchoscopic dilation. (c) A 10-mm x 4-cm nitinol fully covered stent was placed in the trachea to just above the carina.

Recurrent stenosis is an indication for endoluminal stenting or surgery. In our case, tracheal stenosis after bronchoscopic dilation was repeated. Re-dilation with rigid bronchoscopy was performed. Postoperative bilateral pneumothorax, subcutaneous emphysema developed. This time, bronchoscopy revealed partial tracheal laceration in thoracic part of trachea. Tracheal laceration was repaired with right thoracotomy. The inspiratory stridor continued in the patient. A nitinol fully covered stent was placed in the trachea to just above the carina.

With the advancement in the field of thoracic medicine and development of technology, large numbers of patients are now being treated with tracheal stents. Advancement in stent design and development of both covered and uncovered expandable metallic stents have broadened both indications and durability.^[6] Endoluminal stent placement offers a rapid and effective means of opening up and maintaining narrowed airways and result in excellent relief of symptoms and improvement in pulmonary function. However, their use in the pediatric age group is uncommon due to the high incidence of complications, difficult removal and the unclear long-term effect on tracheal growth.^[7] Mostafa and Dessouky reported that endoluminal management of 13 pediatric tracheal stenoses and stenting is a viable option with an acceptable complication rate and minimal effect on tracheal and general growth. Many other studies had an acceptable complication rate with no stent-related mortalities.^[8,9]

Conclusion

It should be noted that ingestion of corrosive agents may cause damage to the upper respiratory tract. Admission for bronchoscopy and esophagoscopy to assess the extent of the injury is warranted in most cases. Tracheal stenosis is the major late complication of caustic airway injury. Careful and accurate stent placement may provide significant and life-saving airway improvement as observed in the presenting case.

Informed Consent: Written, informed consent was obtained from the patient's family for the publication of this case report and the accompanying images.

Peer-review: Internally peer-reviewed.

Authorship Contributions: Concept: H.E.; Design: H.E.; Supervision: H.E.; Resource: H.E., H.D.; Materials: H.E., H.D.; Data: H.E., H.D.; Analysis: H.E.; Literature search: H.E., H.D.; Writing: H.E., H.D.; Critical revision: H.E.

Conflict of Interest: None declared.

Financial Disclosure: The authors declared that this study has received no financial support.

REFERENCES

1. Huang CC, Wu HS, Lee YC. Extensive tracheobronchitis and lung perforation after alkaline caustic aspiration. *Ann Thorac Surg* 2010;89:16703.
2. Represas C, Fernandez A, Leiro V, Botana M, Rodriguez C, et al. Caustic tracheobronchitis. *J Bronchology Interv Pulmonol* 2006;13:156–8.
3. Jaillard S, Nseir S, Métois D, Marquette CH, Darras J, Porte H, et al. Extensive corrosive injuries of the upper airways and gastrointestinal tract. *J Thorac Cardiovasc Surg* 2002;123:186–8. [CrossRef]
4. Mostafa BE, Dessouky OY. The role of endoluminal self-expanding stents in the management of pediatric tracheal stenosis. *Int J Pediatr Otorhinolaryngol* 2008;72:1371–6. [CrossRef]
5. Stephens KE Jr, Wood DE. Bronchoscopic management of central airway obstruction. *J Thorac Cardiovasc Surg* 2000;119:289–96. [CrossRef]
6. Bansal S, Dhingra S, Ghai B, Gupta AK. Metallic stents for proximal tracheal stenosis: is it worth the risk?. *Case Rep Otolaryngol* 2012;2012:450304. [CrossRef]
7. Kumar P, Bush AP, Ladas GP, Goldstraw P. Tracheobronchial obstruction in children: experience with endoscopic airway stenting. *Ann Thorac Surg* 2003;75:1579–86. [CrossRef]
8. Arda IS, Boyvat F, Oğün I, Güney LH, Hiçsönmez A. Preliminary experience with tracheal stent application in children with tracheal stenosis. *Eur J Pediatr Surg* 2007;17:241–3.
9. Cabezalí Barbancho D, Pacheco Sánchez JA, López Díaz M, Tejedor Sánchez R, Benavent Gordo M. The role of tracheobronchial stenting in the management of pediatric airway obstruction. [Article in Spanish]. *Cir Pediatr* 2007;20:175–9.

OLGU SUNUMU - ÖZET

Koroziv asit madde içimi sonrası olan trakeal stenozda yaklaşım: Olgu sunumu

Dr. Hıdır Esme, Dr. Hasan Doğan

Sağlık Bilimleri Üniversitesi, Konya Eğitim ve Araştırma Hastanesi, Göğüs Cerrahi Kliniği, Konya

Sindirim sisteminin koroziv madde ile yaralanması iyi bilinen klinik bir tablo olmakla beraber, kritik olarak hayatı tehdit eden hava yolu hasarı açık bir şekilde dökümanite edilmemiştir. Koroziv asit madde içilmesi ile ortaya çıkan trakeal stenoz çok nadirdir. Biz, isimsiz şişede saklanan asidik temizleme ajanının, kaza ile içilmesinden üç hafta sonra acil kliniğine başvuran dört yaşındaki kız çocuğu olgusunu rapor ettik. Rijit bronkoskopi stenoz varlığını gösterdi. Olgu tekrarlayıcı dilatasyon, trakeal yaralanmanın onarılması ve geçici poliüretan ile kaplı nitinol stentin yerleştirilmesiyle tedavi edildi. Sunduğumuz olguda olduğu gibi dikkatli ve doğru stentin yerleştirilmesi, anlamlı ve hayat kurtarıcı hava yolu iyileşmesi sağlayabilir.

Anahtar sözcükler: Koroziv yaralanma; trakeal darlık; yaklaşım.

Ulus Travma Acil Cerrahi Derg 2021;27(3):377-379 doi: 10.14744/tjtes.2020.21805