

Stent thrombosis in acute coronary syndromes: Patient-related factors and operator-related factors

 Martin Kamenik,  Petr Widimský

Cardiocenter, Third Faculty of Medicine, Charles University and University Hospital Kralovske Vinohrady; Prague-Czech Republic

ABSTRACT

Objective: Stent thrombosis (ST) is a common phenomenon in acute coronary syndromes (ACS) when compared to stable coronary artery disease. This study analyzed the patient- and operator-related risk factors of ST in ACS.

Methods: Coronary angiograms of 1738 consecutive ACS patients admitted in a large tertiary center between year 2014 and 2016 were analyzed retrospectively for the presence of ST. The paired angiograms [ST in ACS during and after percutaneous coronary intervention (PCI)] of the patients were analyzed by two independent observers, with focus on lesion characteristics and procedure techniques. Clinical and laboratory data were collected.

Results: Stent thrombosis was found in 29 (1.6%) ACS patients, with a combination of at least one clinical/laboratory risk factor and one lesion/operator risk factor identified in 28 (96%) out of the 29 ACS patients with ST. The following risk factors for ST were found: Renal insufficiency (OR=4.14, $p<0.001$, 95% CI=1.73-9.88), type 2 diabetes (OR=2.21, $p=0.034$, 95% CI=1.06-4.61), excessive alcohol consumption (OR=3.12, $p=0.023$, 95% CI=1.17-8.33), stent implantation for ST-elevation myocardial infarction (STEMI) (OR=2.28, $p=0.029$, 95% CI=1.08-4.81), left main (LM) or left anterior descending artery (LAD) as culprit lesion (OR=2.80, $p=0.010$, CI 95%=1.27-5.95), and absence of antiplatelet therapy prior to ST (OR=3.58, $p=0.002$, 95% CI=1.60-7.96). The following lesion/operator possible risk factors were identified: Bifurcation lesion ($n=7$; 24%), heavy coronary calcifications ($n=13$; 44%), in-stent restenosis with secondary plate rupture ($n=6$, 20%), inappropriate stent size selection ($n=6$, 20%), and errors in periprocedural drug administration ($n=4$, 14%).

Conclusion: ST occurred in 1/62 ACS patients after PCI. A combination of clinical/laboratory and lesion/operator risk factors were present in almost all ACS patients with ST. This finding may support the search for strictly individualized strategies for the treatment of ACS patients with ST after PCI. (*Anatol J Cardiol* 2020; 24: 274-9)

Keywords: stent thrombosis, risk factors, acute coronary syndrome, antithrombotic treatment

Introduction

Stent thrombosis (ST) is the most dreadful complication of percutaneous coronary intervention (PCI) (1-3). Over the last two decades, coronary angioplasty with stent implantation has gained a firm position in treating patients with coronary vascular disease (CVD). PCI is used alongside with cardiovascular surgery, especially in acute coronary syndrome. Although the incidence of ischemic heart disease is on a decline, the absolute number of coronary atherosclerotic patients treated by implantation of stent is on the increase (4). The benefits of the stent implantation used for acute and elective cases of CVD are indisputable (5). Every medical procedure which leaves an artificial device (such as stent implantation) in an organism carries with certain risks (which in this case is stent thrombosis).

Stent thrombosis is the complete occlusion of a coronary vessel in place of previously implanted stent. This situation usually results in ST-elevation myocardial infarction (STEMI) (6, 7) with very poor prognosis. Mortality rate of this event was reported as high as 40% (8). The prevalence of ST in the modern era of drug eluting stent is between 1.1 and 1.9% (9-12). The clinical risk factors for the development of stent thrombosis are well known.

The previous works on ST suggest acute coronary syndrome as a reason for the stent implantation. However, history of some chronic conditions, vascular dysfunction, and progression of the neoatherosclerosis are the most common risk factors for the development of ST (13-20). In this study we analyzed the clinical, laboratory and operator/lesion characteristics of ACS patients with ST in order to determine the possible risk factors for developing ST.

Address for correspondence: Martin Kamenik, MD, Cardiocenter, Third Faculty of Medicine, Charles University and Faculty Hospital Kralovske Vinohrady, Srobarova 50, Prague-Czech Republic
Phone: +420776573185 E-mail: martinkamenik@gmail.com

Accepted Date: 22.05.2020 **Available Online Date:** 23.09.2020

©Copyright 2020 by Turkish Society of Cardiology - Available online at www.anatoljcardiol.com
DOI:10.14744/AnatolJCardiol.2020.69679



Methods

Coronary angiograms of 1738 consecutive patients (admitted in large tertiary center between year 2014 and 2016) with acute coronary syndrome (ACS) including ST-elevation myocardial infarction (STEMI), non-ST-elevation myocardial infarction (non-STEMI) and unstable angina pectoris were analyzed retrospectively for the presence of ST. There were no exclusion criteria. Paired angiograms (First PCI from the time of stent implantation and second PCI at the time of ST) from 29 ACS patients with ST were analyzed by two independent observers, with focus on lesion characteristics and procedure techniques. The stent type, as well as the post-dilatation procedure followed, was based on the discretion of the operator. Clinical and laboratory data collated from a review of patients' charts were included in the analysis. The analyzed patient characteristics were sex, age, presence of diabetes mellitus 2 type, renal insufficiency, arterial hypertension, dyslipidemia, current smoking status, and regular and excessive consumption of alcoholic beverages (defined as more than 1 standard drink per day). Renal insufficiency was defined as the history of chronic kidney disease stage 3. Patients with type 2 diabetes were included, provided that they were on diet or treated orally with anti-diabetic medications or insulin. The analyzed laboratory characteristic (at time the of stent implantation) were hemoglobin level, creatinine level, troponin level and platelet count. Other characteristics studied were reason for stent implantation (initial diagnosis of STEMI), culprit lesion treated with stent in left main (LM) or left anterior descending artery (LAD), and non-usage of anti-aggregation medication before ST. In our study, only ACS patients with angiographic or post-mortem verified occlusions in stented area were included. The acute and late ST were included according to ARC criteria (21). The follow-up mortality data was obtained from the national health insurance registry. The determination of the possible lesion/operator risk factors for ST was done after a detailed review of literature on the topic, and was decided upon by the investigators.

Statistical analysis

The test for normal distribution of data was by Kolmogorov-Smirnov normality test. Data was expressed as the mean±standard deviations. Student's t-test was done to check for differences between two groups of quantitative data. Chi-square or Fisher's exact test was used (where applicable) to check for the differences between groups of categorical data which were expressed as counts and percentages. The importance of risk factors were subjected to univariate logistic regression. P values less than 0.05 were adopted as significant. All statistical analysis was done using IBM SPSS version 23.

Results

ST was found in 29 (1.6%) of 1738 ACS patients screened. The mortality after two years follow-up was 30%. Two thirds of the

ACS patients with ST were men but were non-significantly different from the women. More than half of the analyzed patients were older than 65 years. The other characteristics which did not differ statistically ($p>0.05$) were history of arterial hypertension, dyslipidemia and smoking. Statistical significant differences were found for renal insufficiency (24% vs. 7%, $p<0.001$), type 2 diabetes (48% vs. 29%, $p=0.034$), alcohol consumption (17% vs. 6%, $p=0.023$), STEMI as reason for stent implantation (58% vs. 36%, $p=0.029$), LM or LAD as a culprit lesion (65% vs. 40%, $p=0.010$) and non-usage of anti-aggregation medications before ST (31% vs. 8%, $p=0.002$). There was a statistical significant difference in the absolute platelet count (243 ± 70 vs. 273 ± 75 , $p=0.005$). The laboratory levels of creatinine, troponin and hemoglobin between the two groups were non-significant ($p>0.05$). The summary of these characteristics are shown in Table 1 and 2. The following risk factors for ST were found: renal insufficiency (OR=4.14, $p<0.001$, 95% CI=1.73-9.88), type 2 diabetes (OR=2.21, $p=0.034$, 95% CI=1.06-4.61), excessive alcohol consumption (OR=3.12, $p=0.023$, 95% CI=1.17-8.33), stent implantation for STEMI (OR=2.28, $p=0.029$, 95% CI=1.08-4.81), LM or LAD as culprit lesion (OR=2.80, $p=0.010$, 95% CI=1.27-5.95), and non-usage of anti-aggregation medications before ST (OR=3.58, $p=0.002$, 95% CI=1.60-7.96). Overview of the risk factors is in Table 3. The average length of hospital stay for the group with ST (6.9 ± 2.7 days) was non-significantly ($p=0.20$) different from that of the control group (5.14 ± 3.43 days).

The stents in the ST were 19.07 ± 6.02 mm in length, with the nominal diameter of 3.32 ± 0.44 mm. There was no significant ($p>0.05$) difference between the nominal diameter and the real diameter after stent implantation. The most frequent stent types used were metallic stents (BMS, 27%) and drug-eluting stents of second generation with everolimus (EES, 24%). The various stent types are shown in Table 4. There were no first generation DES or bioresorbable stents with ST in our study. Post-dilatation was used only in two patients (6%) in the ST group. An average of 1.62 ± 0.55 stent was implanted per patient in the ST group, and this was significantly different ($p<0.001$) from that of the control group (1.15 ± 0.11). A suboptimal result of stenting procedure was more often documented for patients in the ST group [3 (10%) vs. 54 (3%), $p=0.02$].

The possible lesion/operator risk factors of ST can be divided into several parts. The lesion related factors were bifurcation lesion ($n=7$; 24%), heavy coronary calcifications ($n=13$; 44%), in-stent restenosis with secondary plate rupture and ST ($n=6$, 20%). Operator related factors were wrong stent size selection (diameter under-sizing or incomplete lesion coverage) ($n=6$, 20%), and errors in periprocedural drug administration ($n=4$, 14%). The patient-related factor errors in the use of anti-aggregation medications was identified as responsible for ST in 3 patients while the other factors were identified in 4 patients. An overview of the lesion/operator risk factors is shown in Figure 1. A combination of at least one clinical/laboratory risk factor and one lesion/operator risk factor was identified in 28 (96%) out of the 29 ACS patients with ST.

Table 1. Basic clinical characteristics

	Without stent thrombosis, n (%)	With stent thrombosis, n (%)	P-value
Male sex	1205 (70)	24 (82)	0.157
Age above 65 years	1013 (59)	17 (58)	0.619
Type 2 diabetes	507 (30)	14 (48)	0.034
Chronic kidney disease	122 (7)	7 (24)	<0.001
Arterial hypertension	1096 (64)	17 (59)	0.540
Dyslipidemia	567 (33)	13 (45)	0.191
Active smoking	878 (51)	13 (45)	0.485
Alcohol abuse	107 (6)	5 (17)	0.023
STEMI as an indication for the initial PCI	654 (38)	17 (58)	0.029
LM or LAD as culprit vessel	698 (41)	19 (65)	0.010
Absence of antiplatelet therapy prior to ST	191 (11)	9 (31)	0.002
Total	1709 (100)	29 (100)	

LM - left main; LAD - left anterior descending artery

Table 2. Basic laboratory characteristics

	Without stent thrombosis (mean±SD)	With stent thrombosis (mean±SD)	P-value
Creatinine on admission (ug/L)	102±73	98±32	0.297
Troponin on admission (mg/L)	764±1501	1080±2232	0.230
Hemoglobin on admission (g/L)	137±18	132±18	0.083
Platelet count on admission (10⁹/L)	243±70	273±75	0.005




Figure 1. The possible lesion/operator risk factors of ST

Discussion

The main findings of this study is as follows: (1) the mortality rate of ST in our study was in consonant with that of other stud-

ies (22-24); (2) We identified possible lesion/operator risk factors, which are imminent threat of ST, in all patients with ST; (3) the risk of ST was much higher when several risk factors were combined. Despite the technology advancements in stent design and improved antiplatelet therapy, there is an increase in mortality rate of ST patients. As the number of patients treated by PCI increases, there is corresponding increase in the absolute count of the patients who are endangered. The risk factors can be divided into several categories which are characteristics of the patient, lesion characteristics and procedure outcome (25). Our results, considering the patient related risk factors, revealed statistical significant occurrence of stent thrombosis among patients with diabetes mellitus and renal impairment. Similar results have been demonstrated in a study conducted by Tada et al. (26) and Iakovou et al. (27). Excessive alcohol consumption is not frequently described as a risk factor in literature. In 2004, a study on alcoholic consumption and restenosis revealed a lower restenosis rate in patients consuming higher amounts of alcohol (28). The difference in the pathophysiology between restenosis and ST substantiates the findings in our study. The lesion characteristics, significant risk for ACS STEMI as a reason for stenting, and culprit lesion in LM or LAD (29) are not less important. The proposed mechanisms for this finding are malapposition,

Table 3. Risk factors for stent thrombosis

Risk factor	Odds ratio	P-value	95% Confidence interval
Male sex	2.75	0.157	0.76-5.29
Age above 65 years	0.83	0.619	0.41-1.70
Type 2 diabetes	2.21	0.034	1.06-4.61
Chronic kidney disease	4.14	0.001	1.73-9.88
Arterial hypertension	0.79	0.540	0.37-1.67
Dyslipidemia	1.64	0.191	0.78-3.43
Active smoking	0.76	0.485	0.37-1.61
Alcohol abuse	3.12	0.023	1.17-8.33
STEMI as an indication for the initial PCI	2.28	0.029	1.08-4.81
Left main or LAD as culprit vessel	2.75	0.01	1.27-5.95
Absence of antiplatelet therapy prior to ST	3.58	0.002	1.60-7.96

LAD - left anterior descending artery; PCI - percutaneous coronary intervention

Table 4. Stent types with ST

Stent type	n (%)
Everolimus eluting stent	7 (24)
Zotarolimus eluting stent	5 (17)
Biolimus eluting stent	5 (17)
Sirolimus eluting stent	3 (10)
Other type drug eluting stent	1 (3)
Bare metal stent	8 (27)

and stent struts crossing an ostial side branch (30). In addition, there were significantly more suboptimal results from index stent implantation, as well as more stents implanted per patient. This suggest that the complexity of the procedure itself plays a crucial role in ST. Similar findings was reported by Lüscher et al. (31). The percentage of patients who were subjected to post-dilatation was only 6% in ST group. The need for post-dilatation to achieve optimal stent diameter was demonstrated in POSTIT trial. Its positive effect on lowering the rate of ST has also been demonstrated. The heavy calcification and bifurcation lesion as a lesion related risk factors have been well documented (32, 33). The German authors have demonstrated that a decreased stent symmetry in calcified lesions predisposes to malposition of the stent (34). This pathologic findings have suggested that the arterial branch points are predisposed to the development of atherosclerotic plaque and thrombus because they are foci with shear stress (35). The next objects of interest are the stents themselves. The diameter of struts, biocompatibility of polymer containing drug which delays the endothelization, thus preventing restenosis, are important technical characteristics which could influence the development of ST. The dynamics of release of this drug is an important variable that affects the outcome of

the procedure significantly (36). In this study, there were only second generation DES and BMS in the non-randomized setting. We didn't analyze the rate of stent thrombosis between the stent types because DES and BMS were used in the ratio of 2:1. The last set of factors which can predispose to ST is the periprocedural technique of implantation, and the medication given to the patient during and after coronary intervention. Inappropriate stent size selection (diameter under-sizing or incomplete lesion coverage) was identified as the third most frequent predisposing factor for ST. Poorly conducted peri- and post procedural medication (mainly wrong dosage of heparin or non-usage of antiplatelet therapy during or few hours after stent implantation) was identified in 14% of patients with ST. Noncompliance in the usage of dual antiplatelet therapy is a well-known risk factor for ST. The absence of any antiplatelet therapy (ASA or P2Y12 inhibitors) predisposes to statistically higher rate of ST (37). In this study, the risk of ST after discontinuing the antiplatelet therapy was almost four times higher. The importance of patient compliance is therefore imminent. Finally, the combination of risk factors rather than only one, may probably be responsible for the complex events seen in the ST patients in our study. A combination of at least one clinical/laboratory and one lesion/operator risk factor was identified in 28 (96%) out of the 29 patients with ST. This finding suggests that the development of ST is influenced by multiple factors on the side of the patient as well on the side of the operator.

Study limitations

The observational study is by nature selectively bias. The stent type and length were in a non-randomized setting, and was solely the choice of the interventional cardiologist. We did not conduct any advanced visualization for the lesions. On the other hand, the visualization of every implanted stent by IVUS or OCT is inapplicable to conventional or real-life clinical practice, and not

even for patients who are unstable during procedure. Therefore, the review of angiograms was often only guided during by the interventional cardiologist.

Conclusion

Stent thrombosis occurred in 1/62 ACS patients after PCI. A combination of clinical/laboratory and lesion/operator risk factors were present in almost all ACS patients with ST. This finding may support the search for strictly individualized strategies for the treatment of ACS patients with ST after PCI.

Conflict of interest: None declared.

Peer-review: Externally peer-reviewed.

Authorship contributions: Concept – M.K., P.W.; Design – M.K., P.W.; Supervision – M.K., P.W.; Fundings – M.K., P.W.; Materials – None; Data collection and/or processing – M.K.; Analysis and/or interpretation – M.K., P.W.; Literature search – M.K.; Writing – M.K., P.W.; Critical review – M.K., P.W.

References

1. Cutlip DE, Baim DS, Ho KK, Popma JJ, Lansky AJ, Cohen DJ, et al. Stent thrombosis in the modern era: a pooled analysis of multicenter coronary stent clinical trials. *Circulation* 2001; 103: 1967-71.
2. Holmes DR Jr, Kereiakes DJ, Garg S, Serruys PW, Dehmer GJ, Ellis SG, et al. Stent thrombosis. *J Am Coll Cardiol* 2010; 56: 1357-65.
3. Claessen BE, Henriques JP, Jaffer FA, Mehran R, Piek JJ, Dangas GD. Stent thrombosis: a clinical perspective. *JACC Cardiovasc Interv* 2014; 7: 1081-92. [CrossRef]
4. Aggarwal A, Srivastava S, Velmurugan M. Newer perspectives of coronary artery disease in young. *World J Cardiol* 2016; 8: 728-34.
5. Sigwart U, Puel J, Mirkovitch V, Joffe F, Kappenberger L. Intravascular stents to prevent occlusion and restenosis after transluminal angioplasty. *N Engl J Med* 1987; 316: 701-6. [CrossRef]
6. Seto AH, Kern MJ. Early stent thrombosis: Nearly gone, but never forgotten. *Catheter Cardiovasc Interv* 2018; 91: 849-50. [CrossRef]
7. Ergelen M, Gorgulu S, Uyarel H, Norgaz T, Aksu H, Ayhan E, et al. The outcome of primary percutaneous coronary intervention for stent thrombosis causing ST-elevation myocardial infarction. *Am Heart J* 2010; 159: 672-6. [CrossRef]
8. Ge J, Yu H, Li J. Acute Coronary Stent Thrombosis in Modern Era: Etiology, Treatment, and Prognosis. *Cardiology* 2017; 137: 246-55.
9. Inglessis I, Elmariah S, Rengifo-Moreno PA, Margey R, O'Callaghan C, Cruz-Gonzalez I, et al. Long-term experience and outcomes with transcatheter closure of patent foramen ovale. *JACC Cardiovasc Interv* 2013; 6: 1176-83. [CrossRef]
10. Kukreja N, Onuma Y, Garcia-Garcia HM, Daemen J, van Domburg R, Serruys PW; Interventional Cardiologists of the Thoraxcenter (2000 to 2005). The risk of stent thrombosis in patients with acute coronary syndromes treated with bare-metal and drug-eluting stents. *JACC Cardiovasc Interv* 2009; 2: 534-41. [CrossRef]
11. Ergelen M, Uyarel H, Osmonov D, Ayhan E, Akkaya E, Soyulu O, et al. Early stent thrombosis in patients undergoing primary coronary stenting for acute myocardial infarction: incidence, a simple risk score, and prognosis. *Clin Appl Thromb Hemost* 2010; 16: 33-41. [CrossRef]
12. Daemen J, Wenaweser P, Tsuchida K, Abrecht L, Vaina S, Morger C, et al. Early and late coronary stent thrombosis of sirolimus-eluting and paclitaxel-eluting stents in routine clinical practice: data from a large two-institutional cohort study. *Lancet* 2007; 369: 667-78. [CrossRef]
13. Godschalk TC, Gimbel ME, Nolet WW, van Kessel DJ, Amoroso G, Dewilde WJ, et al. A clinical risk score to identify patients at high risk of very late stent thrombosis. *J Interv Cardiol* 2018; 31: 159-69.
14. Mrdovic I, Savic L, Lasica R, Krljanac G, Asanin M, Brdar N, et al. Usefulness of the RISK-PCI score to predict stent thrombosis in patients treated with primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: a substudy of the RISK-PCI trial. *Heart Vessels* 2013; 28: 424-33. [CrossRef]
15. Yadav M, G n reux P, Palmerini T, Caixeta A, Madhavan MV, Xu K, et al. SYNTAX score and the risk of stent thrombosis after percutaneous coronary intervention in patients with non-ST-segment elevation acute coronary syndromes: an ACUITY trial substudy. *Catheter Cardiovasc Interv* 2015; 85: 1-10. [CrossRef]
16. Loh JP, Pendyala LK, Kitabata H, Torguson R, Omar A, Minha S, et al. Comparison of outcomes after percutaneous coronary intervention among different coronary subsets (stable and unstable angina pectoris and ST-segment and non-ST-segment myocardial infarction). *Am J Cardiol* 2014; 113: 1794-801. [CrossRef]
17. Hokimoto S, Tabata N, Yamanaga K, Sueta D, Akasaka T, Tsujita K, et al. Prevalence of coronary macro- and micro-vascular dysfunctions after drug-eluting stent implantation without in-stent restenosis. *Int J Cardiol* 2016; 222: 185-94. [CrossRef]
18. Borovac JA, D'Amario D, Vergallo R, Porto I, Bisignani A, Galli M, et al. Neoatherosclerosis after drug-eluting stent implantation: a novel clinical and therapeutic challenge. *Eur Heart J Cardiovasc Pharmacother* 2019; 5: 105-16. [CrossRef]
19. Joner M, Finn AV, Farb A, Mont EK, Kolodgie FD, Ladich E, et al. Pathology of drug-eluting stents in humans: delayed healing and late thrombotic risk. *J Am Coll Cardiol* 2006; 48: 193-202. [CrossRef]
20. Guo N, Maehara A, Mintz GS, He Y, Xu K, Wu X, et al. Incidence, mechanisms, predictors, and clinical impact of acute and late stent malapposition after primary intervention in patients with acute myocardial infarction: an intravascular ultrasound substudy of the Harmonizing Outcomes with Revascularization and Stents in Acute Myocardial Infarction (HORIZONS-AMI) trial. *Circulation* 2010; 122: 1077-84. [CrossRef]
21. Cutlip DE, Windecker S, Mehran R, Boam A, Cohen DJ, van Es GA, et al.; Academic Research Consortium. Clinical end points in coronary stent trials: a case for standardized definitions. *Circulation* 2007; 115: 2344-51. [CrossRef]
22. Clemmensen P, Wiberg S, Van't Hof A, Deliargyris EN, Coste P, Ten Berg J, et al. Acute stent thrombosis after primary percutaneous coronary intervention: insights from the EUROMAX trial (European Ambulance Acute Coronary Syndrome Angiography). *JACC Cardiovasc Interv* 2015; 8(1 Pt B): 214-20. [CrossRef]
23. Heestermaans AA, van Werkum JW, Zwart B, van der Heyden JA, Kelder JC, Breet NJ, et al. Acute and subacute stent thrombosis after primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: incidence, predictors and clinical outcome. *J Thromb Haemost* 2010; 8: 2385-93. [CrossRef]
24. D'Ascenzo F, Bollati M, Clementi F, Castagno D, Lagerqvist B, de la Torre Hernandez JM, et al. Incidence and predictors of coronary

- stent thrombosis: evidence from an international collaborative meta-analysis including 30 studies, 221,066 patients, and 4276 thromboses. *Int J Cardiol* 2013; 167: 575-84. [\[CrossRef\]](#)
25. Lim S, Koh YS, Kim PJ, Kim HY, Park CS, Lee JM, et al. Incidence, Implications, and Predictors of Stent Thrombosis in Acute Myocardial Infarction. *Am J Cardiol* 2016; 117: 1562-8. [\[CrossRef\]](#)
 26. Tada T, Byrne RA, Simunovic I, King LA, Cassese S, Joner M, et al. Risk of stent thrombosis among bare-metal stents, first-generation drug-eluting stents, and second-generation drug-eluting stents: results from a registry of 18,334 patients. *JACC Cardiovasc Interv* 2013; 6: 1267-74. [\[CrossRef\]](#)
 27. Iakovou I, Schmidt T, Bonizzoni E, Ge L, Sangiorgi GM, Stankovic G, et al. Incidence, predictors, and outcome of thrombosis after successful implantation of drug-eluting stents. *JAMA* 2005; 293: 2126-30. [\[CrossRef\]](#)
 28. Niroomand F, Hauer O, Tiefenbacher CP, Katus HA, Kuebler W. Influence of alcohol consumption on restenosis rate after percutaneous transluminal coronary angioplasty and stent implantation. *Heart* 2004; 90: 1189-93. [\[CrossRef\]](#)
 29. Kuramitsu S, Ohya M, Shinozaki T, Otake H, Horie K, Kawamoto H, et al. Risk Factors and Long-Term Clinical Outcomes of Second-Generation Drug-Eluting Stent Thrombosis. *Circ Cardiovasc Interv* 2019; 12: e007822. [\[CrossRef\]](#)
 30. Mori H, Torii S, Harari E, Jinnouchi H, Brauman R, Smith S, et al. Pathological mechanisms of left main stent failure. *Int J Cardiol* 2018; 263: 9-16. [\[CrossRef\]](#)
 31. Lüscher TF, Steffel J, Eberli FR, Joner M, Nakazawa G, Tanner FC, et al. Drug-eluting stent and coronary thrombosis: biological mechanisms and clinical implications. *Circulation* 2007; 115: 1051-8.
 32. Tasal A, Bacaksiz A, Vatankulu MA, Turfan M, Erdogan E, Sonmez O, et al. Is postdilatation with a noncompliant balloon necessary after coronary stent deployment during primary angioplasty? *J Interv Cardiol* 2013; 26: 325-31. [\[CrossRef\]](#)
 33. Brodie BR, Cooper C, Jones M, Fitzgerald P, Cummins F; Postdilatation Clinical Comparative Study (POSTIT) Investigators. Is adjunctive balloon postdilatation necessary after coronary stent deployment? Final results from the POSTIT trial. *Catheter Cardiovasc Interv* 2003; 59: 184-92. [\[CrossRef\]](#)
 34. Henneke KH, Regar E, König A, Werner F, Klaus V, Metz J, et al. Impact of target lesion calcification on coronary stent expansion after rotational atherectomy. *Am Heart J* 1999; 137: 93-9. [\[CrossRef\]](#)
 35. Iakovou I, Kadota K, Papamantzelopoulos S, Pavlides G, Mitsudo K. Is there a higher risk of stent thrombosis in bifurcation lesion or is it related to the technique? *EuroIntervention* 2010; 6 Suppl J: J107-11.
 36. Torrado J, Buckley L, Durán A, Trujillo P, Toldo S, Valle Raleigh J, et al. Restenosis, Stent Thrombosis, and Bleeding Complications: Navigating Between Scylla and Charybdis. *J Am Coll Cardiol* 2018; 71: 1676-95. [\[CrossRef\]](#)
 37. Sorrentino S, Giustino G, Baber U, Sartori S, Cohen DJ, Henry TD, et al. Dual Antiplatelet Therapy Cessation and Adverse Events After Drug-Eluting Stent Implantation in Patients at High Risk for Atherothrombosis (from the PARIS Registry). *Am J Cardiol* 2018; 122: 1638-46.