



Interobserver and Intraobserver Agreements of the Detection of *Demodex* Infestation by *in Vivo* Confocal Microscopy

Ayse Yildiz Tas,¹ Burak Mergen,² Erdost Yildiz,³ Betul N Bayraktutar,⁴ Ekrem Celik,⁵
 Afsun Sahin,¹ Ceyhun Arici⁶

¹Department of Ophthalmology, Koç University Medical School, Istanbul, Türkiye

²Department of Ophthalmology, University of Health Sciences, Basaksehir Cam and Sakura City Hospital, Istanbul, Türkiye

³Research Center for Translational Medicine, Koç University, Istanbul, Türkiye

⁴Department of Ophthalmology, Tufts Medical Center, Tufts University Faculty of Medicine, Boston, MA, USA

⁵Department of Ophthalmology, Faculty of Medicine, Tekirdağ Namık Kemal University, Tekirdag, Türkiye

⁶Department of Ophthalmology, Cerrahpasa Medical Faculty, Istanbul University-Cerrahpasa, Istanbul, Türkiye

Abstract

Objectives: The purpose of the study was to determine interobserver and intraobserver agreement, repeatability, and intrasubject variation of the detection of *Demodex* infestation in eyelids of blepharitis patients using *in vivo* confocal microscopy (IVCM).

Methods: Eighty-three eyes of 42 blepharitis patients were included in the study. All eyelids were evaluated from temporal to nasal with IVCM using section mode and 10 lashes with their follicles were imagined, and every image with suspicion of *Demodex* infestation was recorded. Two experienced and two inexperienced ophthalmologists were masked for the diagnosis and interpreted the IVCM images regarding the presence of *Demodex* infestation with a 3-week interval. Interobserver and intraobserver agreements were calculated with Cohen's kappa and its variant statistics between and within experienced observers and between inexperienced observers.

Results: While average sensitivity for the diagnosis of demodicosis in IVCM images was 83.35% for experienced and 51.35% for inexperienced observers, the average positive predictive value was 88.6% for experienced observers and 91.05% for inexperienced ones. Interobserver agreement between experienced observers was moderate ($\kappa = 0.529$) and intraobserver agreements within experienced observers were perfect ($\kappa = 0.918$ for observer-1; $\kappa = 0.958$ for observer-2). Interobserver agreement between inexperienced observers was poor ($\kappa = 0.162$) and intraobserver agreements within inexperienced observers were fair ($\kappa = 0.427$ for observer-3; $\kappa = 0.475$ for observer-4).

Conclusion: The sensitivity, interobserver and intraobserver agreements in IVCM image analysis for the detection of *Demodex* infestation were highly associated with the clinical experience on IVCM imaging. In the hands of an experienced clinician, IVCM could be reliable for the diagnosis of ocular demodicosis.

Keywords: Blepharitis, demodex, *in vivo* confocal microscopy, interobserver agreement, repeatability

How to cite this article: Yildiz Tas A, Mergen B, Yildiz E, Bayraktutar BN, Celik E, Sahin A, et al. Interobserver and Intraobserver Agreements of the Detection of *Demodex* Infestation by *in Vivo* Confocal Microscopy. *Beyoglu Eye J* 2022; 7(3): 173-180.

Address for correspondence: Ceyhun Arici, MD. Department of Ophthalmology, Cerrahpasa Medical Faculty, Istanbul University-Cerrahpasa, Istanbul, Türkiye

Phone: +90 532 340 93 51 **E-mail:** ceyhundr@gmail.com

Submitted Date: March 10, 2022 **Accepted Date:** June 23, 2022 **Available Online Date:** August 05, 2022

©Copyright 2022 by Beyoglu Eye Training and Research Hospital - Available online at www.beyoglu-eye.com

OPEN ACCESS This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.



Introduction

Blepharitis is the chronic eyelid margin inflammation and one of the most common conditions encountered in ophthalmology clinics. Blepharitis could affect any age and sex group and may result in dry eye disease by damaging the eyelids and the cornea (1). Confusion on the differential diagnosis of blepharitis and its frequent association with dry eye disease complicates to achieve clear information from prevalence studies. A survey of clinicians in the United States showed that 37–47% of adults patients had findings of blepharitis (2). Venturino et al. studied ocular discomfort in 1148 patients and found that 12% of the subjects had anterior blepharitis, 24% had posterior blepharitis, and 21% had dry eye (3). Chronic blepharitis frequently coexists with meibomian gland dysfunction and irreversibly damage to the ocular surface. Mathers et al. observed that 74% of chronic blepharitis patients have meibomian gland dysfunction (4).

The pathophysiology of blepharitis is still unclear, but its association with staphylococcal superinfection is well established (5). In addition to chronic bacterial infection, (6) environmental factors, (7) and systemic diseases, (1) *Demodex* infestation is also involved in the pathogenesis of blepharitis (8). Recently, a meta-analysis based on 13 published case-control series on the association between *Demodex* infestation and blepharitis indicated that *Demodex* mites were related to chronic blepharitis (9).

Demodex, a genus of small parasitic mites affecting mammals, was first identified in 1841 (10). Although it is the most common ectoparasite in humans, it has recently attracted the attention of clinicians, including ophthalmologists, dermatologists, and other specialists (11). *Demodex folliculorum* and *Demodex brevis* are the only two species that affect the human eye (12). *D. folliculorum* has larger body, about 0.3–0.4 mm long, and clusters as groups in the hair follicles (13). *D. brevis* has a much smaller size, about 0.2–0.3 mm long, lives solitarily in the sebaceous gland (8). Consequently, while *D. folliculorum* is associated with anterior blepharitis, *D. brevis* is associated with posterior blepharitis, meibomian gland dysfunction, recurrent chalazia, and refractory keratoconjunctivitis (14). *Demodex* mites occlude the sebaceous gland orifices, which cause irritation, hyperplasia, and hyperkeratinization in the eyelid margins (15). The cytoskeleton of the mites also may act as a foreign body and cause an inflammatory immune response or granulomatous reaction as implicated in chalazia (16). Furthermore, the secretions of mites may act as a vector that carries bacteria such as staphylococci and streptococci (17). In recent studies, interdependency between demodicosis, staphylococci superinfection, and microbial blepharitis has shown in different patient groups (18). Hereby, accurate, reliable, and repro-

ducible demonstration of *Demodex* infestation will be useful for diagnosis and follow-up of chronic blepharitis cases.

The diagnosis of demodicosis is mainly based on clinical evaluation and cylindrical dandruff (CD) has been reported to be a reliable diagnostic marker in microscopic examination (19). However, it is difficult to diagnose *Demodex* infestation, since there is no macroscopic visible finding and light microscopy examination requires eyelash sampling from the patients. Recently, a non-invasive and painless diagnostic technique, laser scanning in vivo confocal microscopy (IVCM) has been used for diagnosis of *Demodex* infestation and found reliable and effective for ophthalmologic examination (20,21). IVCM is a non-invasive imaging modality that allows real-time imaging of corneal, conjunctival, and meibomian glands structure at a cellular level (22–25). IVCM is also commonly used by dermatologists to diagnose *Demodex* skin infection (26). Despite its widespread and frequent usage, subjective interpretation of IVCM images could negatively affect its reliability and reproducibility (27,28). The role of the observer in IVCM image analysis has been discussed previously in various disease groups (22,23,29). IVCM images are generally interpreted by experienced clinicians, however, many ophthalmologists do not have sufficient experience on IVCM image interpretation. It is critical to know the accuracy of the image interpretation by these relatively inexperienced clinicians.

This study aims to evaluate the role of observer experience in the interpretation of IVCM images at the presence of *Demodex* in patients with seborrheic blepharitis.

Methods

Study Subjects

This study included 83 eyes of 42 consecutive seborrheic blepharitis patients with chronic lid margin inflammation and crusting at the base of the eyelashes. This study was conducted according to the Declaration of Helsinki and it was approved by the local ethical committee. For inclusion in the study, the patients signed informed consent forms. Patients between the ages of 18–55 were included and their age and gender of the patients were recorded as the demographics. All patients had a complete ophthalmological evaluation, including visual acuity testing, slit-lamp biomicroscopic examination, IVCM imaging, and *Demodex* examination was performed. The patients with major ocular disorder, systemic disease, a history of ocular surgery or trauma, the use of contact lenses, or medications that affect the eye were excluded from the study.

IVCM Imaging

All subjects underwent IVCM (Heidelberg Retina Tomography 3 Rostock Cornea Module, Heidelberg Engineering

GmbH, Heidelberg, Germany) with a standard procedure by an experienced ophthalmologist (A.S.). A disposable sterile protective cap (Tomo-Cap, Heidelberg Engineering GmbH, Heidelberg, Germany) was attached to the microscope, and Dexpanthenol 5% (Recugel, Bausch and Lomb, NY, US) was applied as a coupling agent between the cap and the lens objective. Before the examination, a local anesthetic (proparacaine hydrochloride 0.5%) drop was applied to both eyes. After the examiner asked the patient to look toward the light source, the center of the Tomo-Cap was touched gently onto the lower eyelid. The lower eyelid of all eyes was evaluated with IVCM (from temporal to nasal) and approximately 10 lashes and their follicles were imagined, and every suspected image was recorded (Fig. 1). All IVCM examinations were performed by an expert examiner (A.S) and images were obtained using the section mode. A large number of photographs were taken individually, from the root to the tip of an eyelash, to evaluate all the angles of the CD and meibomian orifice. The sections that do not have thin hair or dandruff density were selected and the same examiner recorded the three high-quality images for each eye, 249 images total, without descriptive or patient information.

Optical Microscopy Examination

After IVCM imaging, four eyelashes from the lower lid of each eye were epilated for demodicosis diagnosis. Especially, lashes with a significant amount of CD were targeted for epilation. After the epilation of lashes, one drop of 10% potassium hydroxide (KOH) was added on the lashes and a coverslip was placed carefully on them. The samples in KOH were examined using light microscopy at 10× and 40× by an experienced ophthalmologist (C.A.). The number of *Demodex* in four lashes in total was recorded for each eye. The median value (five per eye) of *Demodex* count was assigned as the cutoff value to create subgroups with low (1-5) and high load (six and above) for *Demodex* infestation.

Image Analysis

Four observers who were masked to the clinical data of patients interpreted the IVCM images regarding the presence of *Demodex*. The first two observers were cornea specialists with more than 3 years of experience in IVCM image interpretation (i.e., experienced observers: Observer-1 and observer-2), and the other two observers were the cornea specialists without any previous experience with IVCM imaging (i.e., inexperienced observers: Observer-3 and observer-4). All observers have read eight full-text articles, original




Figure 1. Representative in vivo confocal microscopy images of eyelash follicles of seborrheic blepharitis patients. The scale bar is 100 μ m. (a) Demodex, which is buried close to the hair, causing dandruff, weakening the bottom of the hair follicle, (b) Demodex buried close to the lash bottom, (c) Demodex infestation, showing three opisthosomas lined up next to each other, (d) single larva buried, (e) two Demodex body, and (f) cluster infestation embedded in the skin close the lash root.

research, and review articles, which were published on IVCM imaging on *Demodex* infestation from March 2012 to March 2017, to diagnose the presence of *Demodex* in IVCM images (18,20,26,30-34). After literature review, all IVCM image set was rated by the observers according to the presence of *Demodex*. All observers determined the results independently when interpreting the IVCM images. The interobserver agreement was evaluated between and within observer groups with different experience levels. In addition to the interobserver agreement, to evaluate the intraobserver agreement, the randomly blended IVCM images were reinterpreted by all observers after 3 weeks. For intraclass correlation, subject eyes were scored from 0 to 3, with the number of images detected by *Demodex* from this eye being 0 for no *Demodex*, 1 for 1–5, 2 for 6–10, and 3 if more than 10.

Statistical Analysis

Statistical analysis was performed using SPSS software (version 26, IBM SPSS Statistics, Chicago, Illinois, USA). The Shapiro–Wilk test was performed to confirm the normal distribution of patient demographics. For each observer, sensitivity, specificity, positive and negative predictive values, and false-positive and false-negative rates for the diagnosis of demodicosis were determined. Diagnosis of demodicosis through light microscopy examination of the lashes was accepted as ground truth for binary classification matrix. Cohen's kappa variant statistics applied to assess interobserver and intraobserver reliability (35). Siegel and Castellan's kappa applied for two rater comparisons with bias correction, Fleiss's kappa applied for more than 2 rater comparison. Kappa values for interobserver and intraobserver reliability were interpreted according to Landis and Koch's guideline (36). To determine the effect of *Demodex* load on the diagnosis of observers, logistic regression analysis was performed. Furthermore, intraclass correlation was performed between observers through the number of demodicosis diagnosed images for each eye of the patients. $P < 0.05$ was considered statistically significant.

Results

The mean age of the patients was 26.14 ± 9.59 years, and 21 (50%) patients were women and 21 (50%) were male. The median number of *Demodex* for each eye was 5 and the interquartile range was between 3 and 7. Among all eyes of the blepharitis patients, 8 (9.6%) eyes did not have demodicosis, 43 (51.8%) eyes had low *Demodex* load, and 32 (38.6%) eyes had high *Demodex* load (Table 1).

For the first image analysis, the average sensitivity for detecting *Demodex* in IVCM images was 83.35% for the experienced observers and 51.35% for the inexperienced observers. The average positive predictive value was 88.6% for experienced observers and 91.05% for inexperienced observers. The sensitivity, specificity, positive and negative predictive values, false-positive and false-negative rates for the first image analysis of each observer are shown in (Table 2).

While interobserver agreement between experienced observers was moderate ($\kappa = 0.529$, $p < 0.001$), interobserver agreement between inexperienced observers was slight ($\kappa = 0.162$, $p = 0.004$). Average Cohen's kappa for interobserver agreements between experienced and inexperienced ob-

Table 1. The subject characteristics

Subjects	Values	
Women (%)	21 (50.0)	
Age (mean \pm SD, year)	26.14 \pm 9.59	
Demodex count (median [IQR])	5.00 [3.00–7.00]	
Presence of Demodex (%)	OD	OS
None	4 (9.5)	4 (9.8)
Low load (1–5)	20 (47.6)	23 (56.1)
High load (>5)	18 (42.9)	14 (34.1)

SD: Standard deviation; IQR: Interquartile range; OD: Oculus dexter; OS: Oculus sinister.

Table 2. The sensitivity, specificity, positive predictive value, negative predictive value, false-positive rate, and false-negative rate of IVCM imaging for the detection of *Demodex* infestation for both experienced and inexperienced observers

Measurement	Experienced observers		Inexperienced observers	
	Observer-1 (%)	Observer-2 (%)	Observer-3 (%)	Observer-4 (%)
Sensitivity	76.0	90.7	70.7	32.0
Specificity	0.0	0.0	25.0	75.0
Positive predictive value	87.7	89.5	89.8	92.3
Negative predictive value	0.0	0.0	8.3	10.5
False-positive rate	100.0	100.0	75.0	25.0
False-negative rate	24.0	9.3	29.3	68.0

servers was 0.198 ± 0.046 (Table 3). Calculated Fleiss's kappa for overall agreement for all observers was 0.192 (slight intraobserver agreement, $p=0.026$)

When the observers repeated IVCM image analysis for intraobserver reliability, they exhibited similar sensitivity, specificity, positive and negative predictive values, false-positive and false-negative rates with the first analysis (Table 4). Intraobserver agreements for experienced observers were in almost perfect agreements ($\kappa=0.918$, $p<0.001$ for observer-1; $\kappa=0.958$, $p<0.001$ for observer-2), compared to moderate agreements of inexperienced observers ($\kappa=0.427$,

$p<0.001$ for observer-3; $\kappa=0.475$, $p<0.001$ for observer-4).

For intraclass correlation, subject eyes were scored from 0 to 3, which are the count of *Demodex* detected image from this eye. Intraclass correlation measurement was applied for intraobserver reliability, as well as for both first and repeat IVCM image analyses of observers (Table 5). Similar to previous inter-rater reliability calculations, interobserver agreement within experienced observer group was higher (0.788, $p<0.001$ for first analysis; 0.730, $p<0.001$ for repeat analysis) than inexperienced observer group (0.349, $p<0.001$ for the first analysis; 0.469, $p=0.002$ for the repeat analysis). Besides

Table 3. Interobserver agreement for detection of Demodex presence and inter-rater reliability comparison in IVCM images in the experienced and inexperienced observer group

Cohen's kappa (κ) = 0.529		
	Observer-2 (experienced)	
	Demodex (+)	Demodex (-)
Observer-1 (experienced)		
Demodex (+)	107	20
Demodex (κ)	37	80
Cohen's kappa (κ) = 0.162		
	Observer-4 (inexperienced)	
	Demodex (+)	Demodex (-)
Observer-3 (inexperienced)		
Demodex (+)	30	71
Demodex (-)	21	122
Cohen's kappa (κ)		
	Inexperienced observers	
	Observer-3	Observer-4
Experienced observers		
Observer-1	0.219	0.194
Observer-2	0.245	0.137

Table 4. Intraobserver reliability, sensitivity, specificity, positive predictive value, negative predictive value, false-positive rate, and false-negative rate of repeated IVCM image analysis for the detection of Demodex infestation for both experienced and inexperienced observers

Measurement	Experienced observers		Inexperienced observers	
	Observer-1	Observer-2	Observer-3	Observer-4
Intraobserver reliability (κ)	0.918	0.958	0.427	0.475
Sensitivity	72.0	88.0	69.3	26.7
Specificity	0.0	0.0	0.0	50.0
Positive predictive value	87.1	89.2	86.7	83.3
Negative predictive value	0.0	0.0	0.0	6.8
False-positive rate	100.0	100.0	100.0	50.0
False-negative rate	28.0	12.0	30.7	73.3

Table 5. Intraclass correlation levels in number of demodicosis diagnosis images per patient between experienced and inexperienced observers for first and repeated image analysis results

Intraclass correlation levels	Experienced Observers		Inexperienced observers	
	Observer-1	Observer-2	Observer-3	Observer-4
Experienced observers				
Observer-1	0.974*	0.788†	0.342†	0.279†
Observer-2	0.730§	0.982*	0.524†	0.286†
Inexperienced observers				
Observer-3	0.499§	0.542§	0.656*	0.349†
Observer-4	0.476§	0.463§	0.469§	0.740*

their higher interobserver agreement levels, experienced observers achieved higher intraobserver agreement levels (0.974, $p<0.001$ for observer-1; 0.982, $p<0.001$ for observer-2) compared to inexperienced observers (0.656, $p<0.001$ for observer-3; 0.740, $p<0.001$ for observer-4) in intraclass correlation. The logistic regression analysis demonstrated that *Demodex* loads of the patients did not affect the sensitivity of IVCM image analysis for *Demodex* detection ($p=0.165$).

Discussion

In our study, we evaluated the sensitivity, specificity, positive and negative predictive values, and inter-rater reliability and the interclass correlation between observers on the diagnosis of *Demodex* infestation through IVCM images from seborrheic blepharitis patients. In our patient sample population for seborrheic blepharitis, 38 (90.5%) patients had visible *Demodex* infestation in eyelash follicles according to the light microscopy examination results. Experienced observers diagnosed these patients with demodicosis with high sensitivity (83.35%). However, for inexperienced observers, the average diagnosis sensitivity was significantly lower (51.35%) for the diagnosis of demodicosis. On the other hand, positive predictive values for all observers were at similar levels regardless of experience. The sensitivity for the diagnosis of *Demodex* with IVCM increased with clinical experience on IVCM. Our result also supported the results of the study by Wang et al. on the interobserver reliability and validity of IVCM imaging for ocular demodicosis (37). While positive predictive value was not affected by the experience of observer, the sensitivity for *Demodex* diagnosis increased with the years of experience.

While the interobserver agreement between experienced observers was moderate ($\kappa=0.529$), both interobserver agreements within the inexperienced group and between experienced and inexperienced observers were poor. Correlatively, intraobserver agreement within experienced ob-

servers was far higher than inexperienced observers. Consistency on the diagnosis of *Demodex* is correlated with the IVCM experience level of the observer. Finally, we validated interobserver agreement results with intraclass correlation levels of observers and we have found a very similar pattern on intraclass correlation levels. Experienced observers achieved significantly higher intraclass correlation levels compared to inexperienced observers. All these interobserver and intraobserver comparisons of *Demodex* diagnosis results show, precise and consistent IVCM image analysis for ocular demodicosis may require a long period of clinical experience with IVCM.

One of the limitations of our study is the miscalculation of specificity due to the small number of seborrheic blepharitis patients without *Demodex* infestation. To pass this limitation, we have also measured the intraclass correlation between observers and we found similar results with inter-rater reliability tests. Another limitation of the study is the non-standardized diagnostic threshold for ocular demodicosis in light microscopy. While Filho et al. (37) suggested the presence of more than 1 *Demodex* in three eyelashes of each eyelid as positive, Wesolowska offered one *Demodex* infected eyelash from 10 random eyelashes was enough for diagnosis (38). We have selected the second one for our diagnostic criteria in light microscopy examination. Furthermore, the overdiagnosis of *Demodex* infestation in the patients by experienced observers leads significant decrease in the specificity of the diagnosis. The main reason for the overdiagnosis could be the increased number of keratinized substances, lipid metabolites, and eyelash scales, which are confused with *Demodex* parts in IVCM imaging.

Blepharitis is a disease highly associated with *Demodex*. The previous studies have shown the prevalence of ocular demodicosis in blepharitis patients between 67.2% and 83.7% (39-42). *Demodex* is not just responsible for blepharitis, it has also significant roles in meibomian gland disorder,

(43) chalazia, (44) pterygium, (45) and rosacea (46). All these diseases are frequently seen in the community and require continuous follow-up for treatment. Although the relatively expensive disposable Tomo-Caps constitute a disadvantage in terms of wide use by ophthalmologists, IVCM is a very convenient option for screening, diagnosis, and follow-up of ocular demodicosis and its related diseases, because of its non-invasive, reliable, and reproducible nature (47).

Conclusion

As a result, even in different experience levels of observers, IVCM showed high sensitivity, positive predictive value, and reproducibility for the diagnosis of ocular demodicosis and IVCM could be safely used for the diagnosis of *Demodex* infestation in ophthalmology clinics.

Disclosures

Acknowledgments: The authors thank to KOÇ University Research Center for Translational Medicine, funded by the Republic of Türkiye Ministry of Development, for allowing us to use their services and facilities. The content of the manuscript is solely the responsibility of the authors and does not necessarily represent the official views of the Ministry of Development, Republic of Türkiye.

Peer-review: Externally peer-reviewed.

Conflict of Interest: None declared.

Authorship Contributions: Concept – A.Y.T., B.M.; Design – E.Y., E.C.; Supervision – A.S., C.A.; Resource – B.M., A.Y.T.; Materials – C.A., B.M.; Data collection and/or processing – B.N.B., E.Y.; Analysis and/or interpretation – B.M., E.Y.; Literature search – A.Y.T.; Writing – E.Ç., E.Y.; Critical review – C.A.

References

- Bernardes TF, Bonfioli AA. Blepharitis. *Semin Ophthalmol* 2010;25:79–83. [CrossRef]
- Lemp MA, Nichols KK. Blepharitis in the United States 2009: A survey-based perspective on prevalence and treatment. *Ocul Surf* 2009;7:S1–14. [CrossRef]
- Venturino G, Bricola G, Bagnis A, Traverso CE. Chronic blepharitis: Treatment patterns and prevalence. *Invest Ophthalmol Vis Sci* 2003;44:774–9.
- Mathers WD, Shields WJ, Sachdev MS, Petroll WM, Jester JV. Meibomian gland dysfunction in chronic blepharitis. *Cornea* 1991;10:277–85. [CrossRef]
- Dougherty JM, McCulley JP. Comparative bacteriology of chronic blepharitis. *Br J Ophthalmol* 1984;68:524–8. [CrossRef]
- O'Brien TP. The role of bacteria in blepharitis. *Ocul Surf* 2009;7:S21–2. [CrossRef]
- Malerbi FK, Martins LC, Saldiva PH, Braga AL. Ambient levels of air pollution induce clinical worsening of blepharitis. *Environ Res* 2012;112:199–203. [CrossRef]
- Luo X, Li J, Chen C, Tseng S, Liang L. Ocular demodicosis as a potential cause of ocular surface inflammation. *Cornea* 2017;36:S9–14. [CrossRef]
- Zhao YE, Wu LP, Hu L, Xu JR. Association of blepharitis with demodex: A meta-analysis. *Ophthalmic Epidemiol* 2012;19:95–102. [CrossRef]
- Desch CE. Human hair follicle mites and forensic acarology. *Exp Appl Acarol* 2009;49:143–6. [CrossRef]
- Elston CA, Elston DM. Demodex mites. *Clin Dermatol* 2014;32:739–43. [CrossRef]
- Nicholls SG, Oakley CL, Tan A, Vote BJ. Demodex species in human ocular disease: New clinicopathological aspects. *Int Ophthalmol* 2017;37:303–12. [CrossRef]
- Lee SH, Chun YS, Kim JH, Kim ES, Kim JC. The relationship between demodex and ocular discomfort. *Invest Ophthalmol Vis Sci* 2010;51:2906–11. [CrossRef]
- English FP, Nutting WB. Demodicosis of ophthalmic concern. *Am J Ophthalmol* 1981;91:362–72. [CrossRef]
- Liu J, Sheha H, Tseng SC. Pathogenic role of demodex mites in blepharitis. *Curr Opin Allergy Clin Immunol* 2010;10:505–10.
- Georgala S, Katoulis AC, Kylafis GD, Koumantaki-Mathioudaki E, Georgala C, Aroni K. Increased density of Demodex folliculorum and evidence of delayed hypersensitivity reaction in subjects with papulopustular rosacea. *J Eur Acad Dermatol Venereol* 2001;15:441–4. [CrossRef]
- Clifford CW, Fulk GW. Association of diabetes, lash loss, and Staphylococcus aureus with infestation of eyelids by Demodex folliculorum (Acari: Demodicidae). *J Med Entomol* 1990;27:467–70. [CrossRef]
- Cheng AM, Sheha H, Tseng SC. Recent advances on ocular Demodex infestation. *Curr Opin Ophthalmol* 2015;26:295–300.
- Gao YY, Di Pascuale MA, Li W, Liu DT, Baradaran-Rafii A, Elizondo A, et al. High prevalence of Demodex in eyelashes with cylindrical dandruff. *Invest Ophthalmol Vis Sci* 2005;46:3089–94. [CrossRef]
- Randon M, Liang H, El Hamdaoui M, Tahiri R, Batellier L, Denoyer A, et al. In vivo confocal microscopy as a novel and reliable tool for the diagnosis of Demodex eyelid infestation. *Br J Ophthalmol* 2015;99:336–41. [CrossRef]
- Wang YJ, Ke M, Chen XM. Prospective study of the diagnostic accuracy of the in vivo laser scanning confocal microscopy for ocular demodicosis. *Am J Ophthalmol* 2019;203:46–52. [CrossRef]
- Efron N, Edwards K, Roper N, Pritchard N, Sampson GP, Shahidi AM, et al. Repeatability of measuring corneal subbasal nerve fiber length in individuals with Type 2 diabetes. *Eye Contact Lens* 2010;36:245–8. [CrossRef]
- Kalteniece A, Ferdousi M, Adam S, Schofield J, Azmi S, Petropoulos I, et al. Corneal confocal microscopy is a rapid reproducible ophthalmic technique for quantifying corneal nerve abnormalities. *PLoS One* 2017;12:e0183040. [CrossRef]
- Randon M, Aragno V, Abbas R, Liang H, Labbe A, Baudouin C. In vivo confocal microscopy classification in the diagnosis of meibomian gland dysfunction. *Eye (Lond)* 2019;33:754–60.

25. Shen YZ, Xu M, Sun S. In vivo confocal microscopy observation of cell and nerve density in different corneal regions with monocular pterygium. *J Ophthalmol* 2020;2020:6506134. [\[CrossRef\]](#)
26. Longo C, Pellacani G, Ricci C, De Pace B, Argenziano G, Zauladek I. In vivo detection of Demodex folliculorum by means of confocal microscopy. *Br J Dermatol* 2012;166:690–2. [\[CrossRef\]](#)
27. Kheirikhah A, Satitpitakul V, Syed ZA, Müller R, Goyal S, Tu EY, et al. Factors influencing the diagnostic accuracy of laser-scanning in vivo confocal microscopy for acanthamoeba keratitis. *Cornea* 2018;37:818–23. [\[CrossRef\]](#)
28. Kheirikhah A, Syed ZA, Satitpitakul V, Goyal S, Muller R, Tu EY, et al. Sensitivity and specificity of laser-scanning in vivo confocal microscopy for filamentous fungal keratitis: Role of observer experience. *Am J Ophthalmol* 2017;179:81–9. [\[CrossRef\]](#)
29. Pacaud D, Romanchuk KG, Tavakoli M, Gougeon C, Virtanen H, Ferdousi M, et al. The reliability and reproducibility of corneal confocal microscopy in children. *Invest Ophthalmol Vis Sci* 2015;56:5636–40. [\[CrossRef\]](#)
30. Cinotti E, Labeille B, Cambazard F, Perrot JL. Reflectance confocal microscopy in infectious diseases. *G Ital Dermatol Venereol* 2015;150:575–83.
31. Jalbert I, Rejab S. Increased numbers of Demodex in contact lens wearers. *Optom Vis Sci* 2015;92:671–8. [\[CrossRef\]](#)
32. Liang H, Randon M, Michee S, Tahiri R, Labbe A, Baudouin C. In vivo confocal microscopy evaluation of ocular and cutaneous alterations in patients with rosacea. *Br J Ophthalmol* 2017;101:268–74. [\[CrossRef\]](#)
33. Sattler EC, Hoffmann VS, Ruzicka T, Braunmühl TV, Berking C. Reflectance confocal microscopy for monitoring the density of Demodex mites in patients with rosacea before and after treatment. *Br J Dermatol* 2015;173:69–75. [\[CrossRef\]](#)
34. Sattler EC, Maier T, Hoffmann VS, Hegyi J, Ruzicka T, Berking C. Noninvasive in vivo detection and quantification of Demodex mites by confocal laser scanning microscopy. *Br J Dermatol* 2012;167:1042–7. [\[CrossRef\]](#)
35. Hallgren KA. Computing inter-rater reliability for observational data: An overview and tutorial. *Tutor Quant Methods Psychol* 2012;8:23–34. [\[CrossRef\]](#)
36. Altman DG. *Practical Statistics for Medical Research*: London: Chapman and Hall; 2020.
37. Filho PA, Hazarbasanov RM, Grisolia AB, Pazos HB, Kaiserman I, Gomes J. The efficacy of oral ivermectin for the treatment of chronic blepharitis in patients tested positive for Demodex spp. *Br J Ophthalmol* 2011;95:893–5. [\[CrossRef\]](#)
38. Wesolowska M, Knysz B, Reich A, Blazejewska D, Czarnecki M, Gladysz A, et al. Prevalence of Demodex spp. in eyelash follicles in different populations. *Arch Med Sci* 2014;10:319–24. [\[CrossRef\]](#)
39. Kabataş N, Doğan A, Kabataş EU, Acar M, Biçer T, Gürdal C. The effect of demodex infestation on blepharitis and the ocular symptoms. *Eye Contact Lens* 2017;43:64–7. [\[CrossRef\]](#)
40. López-Ponce D, Zuazo F, Cartes C, Salinas-Toro D, Pérez-Valenzuela C, Valenzuela F, et al. High prevalence of Demodex spp. infestation among patients with posterior blepharitis: Correlation with age and cylindrical dandruff. *Arch Soc Esp Oftalmol* 2017;92:412–8. [\[CrossRef\]](#)
41. Zeytun E, Karakurt Y. Prevalence and load of Demodex folliculorum and Demodex brevis (Acari: Demodicidae) in patients with chronic blepharitis in the province of Erzincan, Turkey. *J Med Entomol* 2018;56:2–9. [\[CrossRef\]](#)
42. Zhu M, Cheng C, Yi H, Lin L, Wu K. Quantitative analysis of the bacteria in blepharitis with Demodex infestation. *Front Microbiol* 2018;9:1719. [\[CrossRef\]](#)
43. Cheng S, Zhang M, Chen H, Fan W, Huang Y. The correlation between the microstructure of meibomian glands and ocular Demodex infestation: A retrospective case-control study in a Chinese population. *Medicine (Baltimore)* 2019;98:e15595.
44. Yam JC, Tang BS, Chan TM, Cheng AC. Ocular demodicidosis as a risk factor of adult recurrent chalazion. *Eur J Ophthalmol* 2014;24:159–63. [\[CrossRef\]](#)
45. Tarkowski W, Moneta-Wielgoś J, Młocicki D. Do Demodex mites play a role in pterygium development? *Med Hypotheses* 2017;98:6–10. [\[CrossRef\]](#)
46. Kubanov A, Gallyamova Y, Kravchenko A. Clinical picture, diagnosis and treatment of rosacea, complicated by Demodex mites. *Dermatol Reports* 2019;11:7675. [\[CrossRef\]](#)
47. Pellacani G, Witkowski A, Cesinaro AM, Losi A, Colombo GL, Campagna A, et al. Cost-benefit of reflectance confocal microscopy in the diagnostic performance of melanoma. *J Eur Acad Dermatol Venereol* 2016;30:413–9. [\[CrossRef\]](#)