

COMPARISON OF SERUM MARKERS OF MYOCARDIAL ISCHEMIA IN CORONARY ARTERY BYPASS GRAFTING BY ON-PUMP AND OFF-PUMP TECHNIQUES

R. PARVIZI*
M. RAHBANI-NOBAR*
N. SAMADI*
F. KHATIBI*

SUMMARY: Coronary artery bypass grafting (CABG) is usually performed with cardiopulmonary bypass (CPB). However CPB has been associated with several adverse effects. Recently off-pump CABG technique is offered as an alternative to the standard on-pump technique. The purpose of this study was to evaluate the safety of the techniques by estimating and comparing serum markers of myocardial ischemia; CK, CK-MB, LDH, AST, Cardiac Troponin I and some peri- and postoperative complications after the two types of surgical procedures.

Seventy-three patients (58 males and 15 females) with angiographically defined CAD admitted to the hospital for CABG were selected. The off-pump technique was performed in 38 patients with mean age of 60 ± 10.5 years. In 35 other patients with mean age of 55 ± 10.4 years, the on-pump technique was applied. CK, CK-MB, LDH and AST activities and cardiac Troponin I concentration were measured peri- and postoperatively until 24 hours. Characteristics of the patients and some peri- and postoperative data regarding incidence of complication were also recorded.

The activities of serum cardiac enzymes; CK, CK-MB, LDH and AST after on-pump CABG technique were higher than those of off-pump technique ($p < 0.05$ in all cases). The levels of Troponin I were significantly higher after on-pump CABG throughout the entire measurement period ($p < 0.01$) compared off-pump procedure. In the both groups of the patients no preoperative myocardial infarction was observed. In the off-pump technique procedural time was shorter and transfusion of blood and blood products was less than those of on-pump technique, but no differences were noticed between the other recorded data.

Significantly lower release of the enzymes and Troponin I during operation by off-pump technique suggests that the technique causes less myocardial injury. On the basis of little change in the activities of cardiac enzymes and Troponin I in serum and presented data including a marked decrease in blood transfusion and shorter procedural time it is concluded that off-pump CABG is a safe and effective technique in selected patients with appropriate coronary lesions.

Key Words: On-pump CABG, Off-pump CABG, CK, CK-MB, LDH, AST, Troponin I.

INTRODUCTION

The use of cardiopulmonary bypass (CPB) has been linked to a systemic inflammatory response that

may play a role in undesirable patient outcomes (1). Recently off-pump coronary artery bypass grafting (off-pump CABG) has developed and became noticeably important (2). The technique was offered as an alternative to the standard on-pump technique in selected

*From Department of Cardiac Surgery, Shahid Madani Hospital, Faculty of Medicine, University of Medical Sciences, Tabriz, Iran.

and suitable cases. In contrast to the on-pump CABG procedure, this technique avoids CPB and cardioplegia with global cardiac arrest (3) but there is a potential risk of ischemic myocardial injury resulting from normothermic, metabolically active myocardium during the temporary occlusion of the corresponding coronary arteries (4).

Off-pump CABG procedure needs much patience, skill than conventional CABG. The technique was introduced many years ago, but has been abandoned with the advent of CPB and cardioplegic arrest (5, 6). Off-pump CABG operation enables a better investigation of the role of CPB for myocardial protection during coronary bypass operation. It allows to compare global ischemia reperfusion associated with a CABG procedure using CPB, cardioplegia, and global cardiac arrest versus local ischemia reperfusion due to temporary occlusion of one of the coronary arteries as performed by off-pump CABG.

The aim of this study was to evaluate the safety of the technique by measuring the markers of myocardial ischemia including the activities of serum cardiac enzymes and levels of cardiac specific protein release in the serum after on-pump and off-pump CABG procedures. Myocardial injury occurs in essentially 100% of cases following cardiac bypass surgery. Following the injury activities of cardiac enzymes and level of cardiac specific protein increase in the serum. The amount of increase is generally higher for CK-MB than for Troponin I; it has been suggested that Troponin I may be a better marker for distinguishing myocardial infarction from 'normal' preoperative injury (7, 8). Incidence of peri and postoperative complications in off-pump and on-pump CABG was also estimated and compared with each others.

MATERIAL AND METHODS

Subjects

The study included 73 patients (58 males and 15 females) with a mean age of 58 ± 10.3 years referred to Shahid Madani Hospital, Tabriz-Iran, who were scheduled for surgical revascularization. The off-pump CABG group comprised 38 patients with a mean age of 60 ± 10.5 years with one-vessel (n=5), two-vessels (n=11), three-vessels (n=20) and four-vessels (n=2) coronary artery disease (CAD). The on-pump CABG group included 35 patients with a mean age of 55 ± 10.4 years with one-

vessel (n=3), two vessels (n=6), three-vessels (n=20) and four-vessels (n=6) CAD. Characteristics of the patients and some pre- and postoperative data were recorded. To ensure equivalency between the two groups, patients had to be appropriate candidates for either procedures. Both groups of the patients underwent bypass grafting through a sternotomy incision.

Sampling

The tests were carried out on peripheral venous blood samples collected in simple tubes. The samples were obtained before and 6, 9, 12 and 24 hours after operation.

Measurements

Catalytic activities of CK, LDH and AST in serum samples were determined by using commercial reagents (Boehringer Mannheim, Mannheim, Germany) in an automated chemical analyzer (Cobas Mira) at 25°C and results reported as IU/L. CK-MB isoenzyme activity was detected by immunoinhibition with commercial agents (Boehringer Mannheim) in the same analyzer at 25°C. Cardiac Troponin I concentration in the serum samples were assayed by using Troponin I enzyme immunoassay test kit (Diagnostic Automation Inc. Calabasas, CA 91302 USA) in Awareness stat Fax 2600 analyzer. The ELISA test is based on the principle of solid phase enzyme linked immunosorbent assay.

Statistical analysis of data was accomplished by means of the SPSS 100 statistical software package. Regression analysis was calculated for correlation between parameters. A p value of the less than 0.05 was considered significant. All data are expressed as the mean \pm SD.

RESULTS

There was no hospital mortality, no neurologic accidents, pulmonary insufficiency or incidences of myocardial infarction in the on-pump and off-pump CABG groups. In the off-pump technique procedural time was shorter than that of on-pump CABG technique ($p < 0.05$). Blood transfusion in on-pump CABG group was required more frequently than off-pump CABG group and the differences were marked ($p < 0.05$). Inotrop requirements in both procedures were almost similar ($p > 0.05$).

The cardiac enzymes activities; CK, CK-MB, LDH and AST were measured in the serum samples obtained before and after operation and the results are reported as IU/L. Comparison of the enzymes activities obtained by different procedures are shown in Table 1.

Table 1: The mean±SD values of cardiac enzymes activities in serum before and after operation by off-pump and on-pump procedures at different time points.

Enzymes IU/L	Before operation		After operation					
	Off-pump n=38	On-pump n=35	6 hours		12 hours		24 hours	
			Off-pump n=38	On-pump n=35	Off-pump n=38	On-pump n=35	Off-pump n=38	On-pump n=35
CK*	131±96	122±60	841±84	1324±63	1960±990	2042±885	1851±1020	1917±985
CK-MB*	17±6	20±4	38±16	45±12	31±14	53±17	29±18	38±18
LDH*	413±130	451±111	470±140	666±141	673±220	801±141	693±231	850±205
AST*	26±8	24±6	36±10	55±12	68±17	87±14	73±18	97±12

n: number of patients

*: Significant difference between off-pump and on-pump procedure ($p < 0.05$)

Before operation the activities of CK in both groups were almost similar ($p > 0.05$) but after surgery increased in the both groups of the patients. Peak values for CK were measured at 12 hours. The elevation of the enzyme in on-pump technique was significantly higher than that of off-pump technique in all the postoperative samples ($p < 0.05$ in all cases). The activities of CK-MB isoenzyme before operation were at normal range and increased after operation with a peak at 6 and 12 hr samples in off-pump and on-pump procedures respectively. Meaningful differences were noticed between the two groups ($p < 0.05$). The activities of two other enzymes; LDH and AST were also measured in all samples. After operation the activities of LDH and AST increased and peak of activities appeared at 12 hr samples and remained at that level until 24 hr samples. Activities of both enzymes in on-pump procedure were significantly higher than off-pump procedure ($p < 0.05$ in all cases).

The concentration of Troponin I was measured pre- and postoperatively. As shown in Table 2 before operation the levels of Troponin I in serum was very low and sometime non-detectable, but after operation its concentration increased in both techniques and the maximum of Troponin I concentration was detected at 9 and 12 hr after operation in off-pump and on-pump procedures respectively. Comparing the results of both groups the concentration of Troponin I in all post-operative samples of on-pump technique was higher than that of on-pump procedure ($p < 0.05$).

To increase specificity of studied parameters the relationship between the numbers of grafts and the enzyme activities or protein levels at postoperative samples with maximum activities or levels were estimated. The patients in both groups were subdivided into four subgroups according to the number of grafts received and performing Pearson's correlation coefficient the relationship between the number of the

Table 2: The mean levels of Troponin I in the serum of patients before and after operation by off-pump and on-pump techniques.

Procedures	Mean±SD levels of Troponin I (ng/ml)				
	Before operation	After operation			
		6 hours	9 hours	12 hours	24 hours
Off-pump	0.75±0.28	3.55±0.28	3.34±0.31	3.25±0.54	2.15±0.52
On-pump	0.63±0.33	6.46±2.35	9.21±1.95	6.85±1.55	4.8±2.30

Table 3a: Correlation between the number of vessels grafted and variables; CK, CK-MB, LDH, AST and Troponin I after off-pump operation.

Variables	Number of vessels grafted							
	1		2		3		4	
	r	p	r	p	r	p	r	p
CPK	0.22	0.41	0.380	0.24	0.415	0.33	0.357	0.28
CK-MB	0.44	0.03	0.59	0.019	0.620	0.031	0.389	0.016
LDH	0.197	0.88	0.53	0.29	0.133	0.45	0.221	0.18
AST	0.549	0.43	0.49	0.16	0.312	0.22	0.292	0.32
Troponin I	0.203	0.016	0.33	0.032	0.292	0.009	0.312	0.012

Table 3b: Correlation between the number of vessels grafted and variables; CK, CK-MB, LDH, AST and Troponin I after on-pump operation.

Variables	Number of vessels grafted							
	1		2		3		4	
	r	p	r	p	r	p	r	p
CPK	0.549	0.95	0.434	0.081	0.610	0.73	0.440	0.48
CK-MB	0.221	0.88	0.133	0.089	0.341	0.153	0.770	0.90
LDH	0.070	0.182	0.349	0.331	0.435	0.10	0.125	0.18
AST	0.012	0.67	0.291	0.680	0.380	0.30	0.480	0.075
Troponin I	0.320	0.045	0.144	0.028	0.075	0.004	0.380	0.008

grafts and the variables was assessed. As shown in the Tables 3a and 3b in both techniques the regression analysis demonstrated a positive correlation between the number of vessels grafted and levels of Troponin I, but in the case of CK-MB the correlation was only positive and significant in off-pump group. In spite of higher activities of the other enzymes in the samples obtained after on-pump technique there was no correlation between the number of grafts and the procedures.

DISCUSSION

Open-heart surgery, such as aortic surgery and coronary artery bypass grafting (CABG) is usually accompanied by ischemic or mechanical damage to the heart tissues. Postoperative cardiac failure due to myocardial necrosis still remain the major complication in cardiac surgical procedures and the main cause of increased

morbidity and mortality. Significant myocardial injuries associated with cardiac surgery result in significant (up to 10-fold) increases in the two-year complication rates (10). In recent years new surgical designs in the treatment of coronary artery disease have generated controversy and debate (11,12). Early results suggest that off-pump coronary artery bypass grafting is a safe and effective means of coronary revascularization with excellent short-term results and minimal morbidity (13). However, the safety and efficacy of the procedure are still to be determined. The main object of this prospective study was to compare postoperative kinetic and patterns of cardiac Troponin I and CK, CK-MB, LDH and AST activities after off-pump CABG versus conventional on-pump CABG.

We found no differences in the incidence of postoperative complication between the off-pump CABG and CABG with CPB groups. Because the rates of many

complications in this center are in the range of 1-5 percent to show a significant difference between groups would require a sample size of well over 1000 patients. In this study we also found meaningful differences between two groups with regard to overall length of stay, blood and blood products transfusion and length of procedure. They were all low in off-pump CABG procedure. Similar results have been reported by others (14, 15).

All patients undergoing CABG surgery with or without CPG postoperatively showed an increase of cardiac Troponin I levels. After uncomplicated coronary revascularization, patients with the off-pump CABG technique continuously showed lower serum Troponin I concentrations than those with the on-pump CABG technique. The release following CABG with or without CPG were different; Troponin I reached its postoperative peak value in patient with the off-pump CABG technique earlier than those with the on-pump CABG technique. Different time pattern (peak) and increased levels of the protein after CABG operation have been reported by others (16, 17). The differences may be due to length of operation and method of detection of Troponin I in different laboratories.

The present study reveals that an uncomplicated CABG without signs of preoperative MI causes a moderate increase of the following enzymes: CPK, CK-MB, LDH and AST. Patients with the off-pump CABG presented low activities of the enzymes than patients with the on-pump. The findings are compatible with those of reported by Kilger *et al.* (4). Time patterns of elevation of the enzymes in both operation procedures were the same except in the case of CK-MB that reached to maximum in off-pump technique earlier than on-pump technique (6 hr postoperatively versus 12 hr).

Studying specificity of the markers of myocardial ischemia by performing Pearson's correlation coefficient of the enzyme activities or protein concentrations and number of vessels grafted revealed positive correlations in the case of Troponin I. In this study the increase of the CK-MB and protein, particularly Troponin I which are reliable and highly specific markers of myocardial ischemia during cardiac operation can be used to assess the efficacy of cardio protective procedures (18). It has to be noted that reversible myocardial ischemia can cause a functional dysintegrity of cell membranes and release

of cytosolic enzymes without subsequent cellular necrosis (19). Since no patient developed MI as confirmed by ECG and echocardiography increased activities or levels of protein following open-heart surgery may be due to temporary myocardial ischemia that leads to release of cytosolic enzymes or proteins from reversibly injured cardiac cells.

Significantly greater extant release of the markers in the on-pump CABG procedure may be partially due to reperfusion of heart following global cardiac arrest, oxidative stress and inflammation (20). In view of low activities or level of the marker in the patient undergoing off-pump CABG our data do not support additional release of the markers induced by myocardial injury due to Octopus devices.

Marked correlation between levels of Troponin I and the number of vessels grafted suggests that Troponin I is the best indicator of the myocardial injury. On the basis of lower release of the markers in off-pump-CABG procedure it was concluded that the technique is safe and effective.

REFERENCES

1. Arom KV, Flavin TF, Emery WR, Kshetry VR, Janey PA, Petersen RJ : Safety and efficacy of off-pump coronary artery bypass grafting. *Ann Thorac Surg*, 69:704-710, 2000.
2. Scheld H, Schmid C : Cardiac surgery without the use of cardiopulmonary bypass: The challenges. *Curr Opin Anesth*, 11:5-8, 1998.
3. Borst C, Santamore WP, Smedira NG, Bredee JJ : Minimally invasive coronary artery bypass grafting: on the beating heart via limited access. *Ann Thorac Surg*, 63:1-5, 1997.
4. Kilger E, Pichler B, Weis F, Goetz A, Lamm P, Schutz A, Muchlbayer D, Frey : Markers of myocardial ischemia after minimally invasive and conventional coronary operation. *Ann Thorac Surg*, 70:2023-2028, 2000.
5. Buffolo E, Andrade JCA, Branco JNR, Aguiar LF, Ribeiro EE, Jatene AD : Myocardial revascularization without extra corporeal circulation: Seven-year experience in 593 cases. *Eur J Cardiothorac Surg*, 4:504-509, 1990.
6. Benetti FJ, Naselli G, Wood M, Geffner L : Direct myocardial revascularization without extra corporeal circulation: Experience in 700 patients. *Chest*, 100:312-316, 1991.

7. Dufour DR, Lott JA, Henry JB : *Clinical Enzymology in: Clinical Diagnosis and management by Laboratory methods*. Henry, Twentieth Ed, Saunders, pp 281-303, 2001.
8. Apple F : *Acute myocardial infarction and coronary reperfusion: Serum cardiac markers for the 1990's*. *Am J Clin Path*, 97:217-226, 1992.
9. Engall E : *Methods in Enzymology*. Vol 70, Ed by H Van Vunakis and JJ Langone, Academic Press, New York, pp 419-492, 1980.
10. Haggart PC, Ludman PF, Bradbury AW : *Cardiac Troponin: a new biochemical marker for peri-operative myocardial injury*. *Eur J Vase Endovase Surg*, 22:301-305, 2001.
11. Bonchek LI, Ulliyot DJ : *Minimally invasive coronary bypass: A dissenting opinion*. *Circulation*, 98:495-497, 1998.
12. Mack M, Damiano R, Matheny R, Reichenspurner H, Carpentier A : *Inertia of success: a response to minimally invasive coronary bypass: Dissenting opinion*. *Circulation*, 99:1404-1406, 1999.
13. Turner WF : *"Off-pump" coronary artery bypass grafting: The first one hundred cases of the Rose City experience*. *Ann Thorac Surg*, 68:1482-1485, 1999.
14. Buffolo E, Andrade JCS, Branco JNR, Teles CA, Aguiar LF, Gomes WJ : *Coronary artery bypass grafting without cardiopulmonary bypass*. *Ann Thorac Surg*, 61:63-66, 1996.
15. Ascione R, Lloyd CT, Underwood MJ, Lotto AA, Pitsis AA, Angelini GD : *Economic outcome of off-pump coronary artery bypass surgery: A prospective randomized study*. *Ann Thorac Surg*, 2:216-221, 1999.
16. Mair J, Larue C, Mair P, Balogh D, Calzolari C, Puschendorfer B : *Use of cardiac Troponin I to diagnose preoperatively myocardial infarction in coronary artery bypass grafting*. *Clin Chem*, 40:2066-2070, 1994.
17. Peivandi AA, Hake U, Dahm M, Opfermann UT, Peetz D, Hafner G, Loos AH, Tzanova I, Qelert H : *Coronary revascularization: off-pump versus on-pump a comparison of behavior of biochemical cardiac ischemia markers*. *Z Cardiol*, 91:203-211, 2002.
18. Katrukha A, Troponins AMI : *Diagnostics and what else? Clinical Laboratory International*, 26:14-16, 2002.
19. Piper HM, Schwartz P, Spahr R, Hutter JF, Spieckermann PG : *Early enzyme release from myocardial cell is not due to irreversible cell damage*. *J Mol Cell Cardiol*, 16:385-388, 1984.
20. Matata BM, Sosnowski AW, Galinanes M : *Off-pump bypass graft operation significantly reduces oxidative stress and inflammation*. *Ann Thorac Surg*, 69:785-791, 2000.

Correspondence:
M. Rahbani-Nobar
Dept. of Cardiac Surgery,
Shahid Madani Hospital,
Faculty of Medicine,
University of Medical Sciences,
Tabriz, IRAN.
e-mail: rahbanim@hotmail.com