

Three-Year Assault Cases performed on the Emergency Trauma Center

Acil Travma Merkezine Başvuran Üç Yıllık Darp Olgularının Analizi

© Melih Yucel SANLIER¹, © Behcet AL¹, © Volkan CELEBI², © Gorkem Alper SOLAKOGLU¹, © Eltaf TORUN¹, © Kurtulus ACIKSARI¹, © Bilgehan Ahmet CUMHUR³, © Nafiye SANLIER¹

¹Istanbul Medeniyet University Faculty of Medicine, Department of Emergency, Istanbul, Türkiye

²Serik State Hospital, Clinic of Emergency Service, Antalya, Türkiye

³Viransehir State Hospital, Clinic of Emergency Service, Sanliurfa, Türkiye

ABSTRACT

Objective: This study aims to compare the results of cases with a history of assault presented to our emergency department over a period of three years with literature data and to share our experiences with colleagues.

Methods: Demographic characteristics, trauma findings, clinical conditions, surgical interventions, mortality, and outcomes of patients with a history of assault to the emergency department between 2020 and 2022 were retrospectively examined. The results were analyzed using the Number Cruncher Statistical System 2007 Statistical Software package program (Utah, USA).

Results: A total of 2900 patients, including 1850 males (63.79%) and 1050 females (36.21%), with an average age of 35.53±12.46 years, were included in the study. Most admissions (31.28%) occurred during the summer months and outside working hours. Males inflicted all injuries. The most common injuries occurred in the head-neck region (63.31%) and upper extremities (47.97%) in both sexes. Facial injuries were significantly more common in males ($p<0.05$), whereas spinal and pelvic injuries were common in females ($p<0.05$). Tenderness, edema, hematoma, abrasion, ecchymosis, and soft tissue laceration were the most frequently observed findings. Female patients were more likely to be discharged home ($p=0.0001$), whereas male patients had a higher hospitalization rate ($p=0.0001$). Life-threatening ailments were discovered in 23 patients (0.79%). Surgical intervention was performed in 50 patients (1.72%). No patient experienced fatal outcomes.

Conclusions: All assaults were committed by males. Males are more frequently and severely assaulted than females. The victims were mainly from young people, and assaults predominantly occurred during the summer months and outside working hours. Head-neck and extremity injuries were the most common. There were no fatal outcomes. Nine of the 10 patients were discharged to home from the emergency department.

Keywords: Assault, emergency, sex, education, age

ÖZ

Amaç: Bu çalışmanın amacı acil servisimize üç yıllık süre içinde başvuran darp öyküsü olan olguların sonuçlarını literatür verileri ile karşılaştırmak ve deneyimlerimizi meslektaşlarımızla paylaşmaktır.

Yöntemler: 2020-2022 yılları arasında 3. basamak bir hastane acil servisine saldırı öyküsü ile başvuran olguların demografik özellikleri, travma bulguları, klinik durumları, cerrahi müdahaleleri, mortaliteleri ve hasta sonuçları retrospektif olarak incelendi. Sonuçlar Number Cruncher Statistical System 2007 Statistical Software paket programı (Utah, ABD) kullanılarak analiz edilmiştir.

Bulgular: Yaş ortalaması 35,53±12,46 yıl olan 1850 erkek (%63,79) ve 1050 kadın (%36,21) olmak üzere toplam 2900 hasta çalışmaya dahil edildi. Başvuruların çoğunluğu (%31,28) yaz aylarında ve çalışma saatleri dışında gerçekleşmiştir. Tüm yaralanmalar erkekler tarafından gerçekleştirilmiştir. Her iki cinsiyette de en sık yaralanmalar baş-boyun bölgesinde (%63,31) ve üst ekstremitelerde (%47,97) meydana gelmiştir. Yüz yaralanmaları erkek hastalarda istatistiksel olarak daha sık görülürken ($p<0,05$), omurga ve pelvis yaralanmaları kadın hastalarda daha sıkı ($p<0,05$). Hassasiyet, ödem, hematoma, abrazyon, ekimoz ve yumuşak doku laserasyonu en sık gözlenen bulgulardır. Kadın hastaların eve taburcu edilme olasılığı daha yüksekken ($p=0,0001$), erkek hastaların hastaneye yatış oranı daha yüksekti ($p=0,0001$). Hayati tehlike 23 hastada (%0,79) tespit edildi ve 50 hastaya (%1,72) cerrahi müdahale uygulandı. Hiçbir hastada ölümcül sonuç görüldü.

Sonuçlar: Çalışmamız, tüm saldırı olgularının erkekler tarafından gerçekleştirildiğini ve erkeklerin kadınlara kıyasla daha sık ve ağır saldırıya uğradığını ortaya koymuştur. Mağdurlar çoğunlukla genç yaş grubundadır ve saldırılar ağırlıklı olarak yaz aylarında ve mesai saatleri dışında gerçekleşmiştir. Saldırılarda en sık baş-boyun ve ekstremitelerde yaralanmaları görülmüş, ölümlerle sonuçlanan vaka olmamış ve on hastadan dokuz acil servisten eve taburcu edilmiştir.

Anahtar kelimeler: Saldırı, acil durum, cinsiyet, eğitim, yaş

Address for Correspondence: B. Al, İstanbul Medeniyet University Faculty of Medicine, Department of Emergency, İstanbul, Türkiye

E-mail: behcetal@gmail.com **ORCID ID:** orcid.org/0000-0001-8743-8731

Received: 10 October 2024

Accepted: 14 November 2024

Online First: XXXXX

Cite as: Sanlier MY, Al B, Celebi B, Solakoglu GA, Torun E, Aciksari B, Cumhur BA, Sanlier N. Three-Year Assault Cases performed on the Emergency Trauma Center. Medeni Med J.



INTRODUCTION

The World Health Organization (WHO) defines violence as “events that involve the intentional use of physical force or power, threatened or actual, against oneself, another person, or a group or community, resulting in injury, death, psychological harm, poor development, or deprivation or with a high likelihood of occurrence”. “Assault” is a form of interpersonal violence and ranks as the fourth leading cause of injury-related deaths¹. There was an observed increase in emergency department admissions due to intentional inflicted injuries related to domestic violence, friend violence, and, notably, violence against women. These events not only disrupt victims’ sense of security and quality of life but also lead to psychological and emotional consequences for family members, friends, and the community at large^{2,3}. Materials such as fists, kicks, stones, weapons, and sticks are commonly used as attack tools to harm other objects⁴. Recognizing the significance of injuries in this context is crucial for evidence-based approaches to prevent death and disability and mitigate harm. These cases are assessed as “medico-legal cases” in emergency department admissions.

Globally, the mortality rate of assault cases varies, with males having double the mortality rate of females. One in three women aged above 15 years is reported to have been assaulted by an intimate partner⁵. More than 90% of interpersonal violence cases occur in low-and middle-income countries⁶. Assault cases tend to present more frequently to emergency departments during the summer months and outside working hours⁷. The head-neck regions are most commonly attacked, with bruises, swelling, fractures, contusions, soft tissue lacerations, and bleeding being the most prevalent findings⁸. The severity of an incident may not be directly proportional to the extent of the lesions⁹. Applying the advanced trauma life support algorithm is essential in patient management. Global reactions to such cases have been increasing in recent years, and these actions are raising awareness.

Emergency physicians play a significant role in the clinical evaluation and management of patients with a history of assault. The pattern of presentation of assaulted patients to the emergency department, their physical and psychological conditions, injury localisations and epidemiological characteristics of the victims help to understand and manage such incidents more accurately. The level of education and awareness of healthcare professionals also play a vital role in managing such cases¹⁰. It is of utmost importance for healthcare professionals, especially emergency physicians, who first encounter this patient group clearly and transparently

document injuries and treatment management, which is crucial to ensure that the rights of the parties involved are not violated¹¹. In this study, a retrospective evaluation was performed on the demographic characteristics, presentation patterns, trauma localizations, clinical conditions, treatments, and out-comes of patients with a history of assault admitted to the emergency department of tertiary hospital with a history of assault between 2020-2022. This study aimed to share our clinical experiences with emergency physicians and to raise awareness of assault both clinically and socially.

METHOD

Study and Location

Cases: Presented to the emergency department of tertiary hospital with a “history of assault” were retrospectively examined between January 01, 2020 and December 31, 2022. This study was approved by the H.S. Istanbul Medeniyet University Göztepe Training and Research Hospital Clinical Research Ethics Committee (decision no :2023/0219, date: 29.03.2023). The study was conducted in accordance with the ethical standards of the 1964 Helsinki Declaration (Appendix 3 of the Helsinki Declaration) and its subsequent amendments. The location of the survey, emergency department of tertiary hospital in Istanbul, is among the neighborhoods with the highest educational levels in the metropolitan city.

Data collection: Between the specified dates, cases with ICD codes W50 (assault by another person, striking, stomping, bending, biting) and W51 (assault by another person) were screened. Clinical and forensic case information found in the digital files of the cases were accepted as data. The data were recorded in an Excel file under main headings. After data collection, the data were transferred to an NCSS file for analysis.

Parameters screened: The data recorded for analysis in this study are as follows:

1. Demographic data of patients: age, sex, date, and time of admission
2. Glasgow Coma Scale at the time of admission
3. Life-threatening situation
4. Localization of assault (including sub-regions in the head such as nasal, maxilla, mandible, zygoma, frontal, orbital, temporal, parietal, occipital, neck, and spinal injuries, thoracic injuries, abdominopelvic injuries, extremity injuries)
5. Physical examination findings related to the assault (tenderness, edema, hematoma, abrasion, ecchymosis, presence of cuts, fractures, lacerations, etc.)

6. Imaging results (X-ray, computed tomography)
7. Consultations requested.
8. Treatments performed (surgery, dressing, incision suturing, bone reduction, and splint application, prescription only, no treatment)
9. Outcomes (discharged home, outpatient follow-up, hospitalization, exists, treatment refusal, and unauthorized abandonment).

Inclusion criteria:

1. Patients aged 18 years who presented between January 01, 2020 and December 31, 2022
2. Patients diagnosed with a diagnosis entered digital files with codes W50 and W51.

Exclusion criteria:

1. Patients who were admitted to the emergency department with the request of "being assaulted" but were later identified as "victims of verbal violence only" during the investigation, and cases in which an entry-exit report under police supervision" was requested due to an unspecified crime were excluded from the study.

2. During the three-year data screening phase, 3515 patient files were accessed. Of these, 2900 patients who satisfied the study criteria were recorded.

Statistical Analysis

In this study, statistical analyses were performed using the NCCSS (Number Cruncher Statistical System) 2007 Statistical Software package program (Utah, USA). Descriptive statistical methods (mean, standard deviation) were employed to evaluate the data. Additionally, the Shapiro-Wilk normality test was used to assess the distribution of variables. For variables with a normal distribution, one-way analysis of variance was used for intergroup comparisons, Tukey's multiple comparison test for subgroup comparisons, and the chi-square test for comparisons of qualitative data. The results were evaluated at a significance level of $p < 0.05$.

RESULTS

Among the patients included in our study, 1850 (63.79%) were male and 1050 (36.21%) were female. The mean age of all patients was determined to be 35.53 ± 12.46 (minimum 18, maximum 92) years. The distribution of patients who presented in 2020, 2021, and 2023 was 29.00 ($n=841$), 33.62 ($n=975$), and 37.38 ($n=1084$), respectively.

The highest number of presentations occurred during the summer months (June, July, and August), at 31.28%. More than half of the patients (54.14%) were admitted to the emergency department between 3:00 and midnight.

When the region and localization of the trauma were examined, the head (63.31%) and upper extremities (47.97%) were most commonly injured, respectively. The maxilla (16.79%) within the head region showed the highest injury rate. The recorded patient findings included tenderness, edema, hematoma, abrasion, ecchymosis, and soft tissue laceration. Tenderness was the most frequently observed sign in the trauma region (62.45%). Among the 2900 patients included in the study, 23 (0.79%) were considered life-threatening, and 50 (1.72%) were hospitalized in surgical clinics (Table 1). Patients were predominantly consulted in the plastic surgery (8.31%) and orthopedic (7.48%) departments (Table 1).

Nasal ($p=0.0001$), mandibular ($p=0.045$), frontal ($p=0.0001$), orbital ($p=0.0001$), and temporal ($p=0.025$) trauma occurred at higher rates in male patients than in female patients. The incidence of spinal ($p=0.006$), pelvic ($p=0.0001$), lower extremity ($p=0.0001$), and upper extremity ($p=0.0001$) trauma was higher in female patients (Table 2). Surgical hospitalization and out-patient follow-up were higher in male patients, whereas the discharge rate in female patients was higher ($p=0.0001$) (Table 2). Tenderness ($p=0.001$) and ecchymosis ($p=0.0001$) were more prevalent in female patients, whereas laceration ($p=0.0001$), edema ($p=0.0001$), hematoma ($p=0.007$), and abrasion ($p=0.002$) were more prevalent in male patients (Table 2). Discharge rates were higher among patients arriving during the day ($p=0.0001$), whereas hospitalizations to surgical departments were higher in patients arriving more toward the evening and night ($p=0.02$). The mean age ratio was lower among patients arriving outside working hours ($p=0.0001$) (Table 3).

Patients who refused treatment and left the emergency department had a higher incidence of nasal ($p=0.0001$) and cervical ($p=0.011$) injuries. Patients with lower extremity and orbital injuries were more likely to be hospitalized and were treated ($p=0.0001$) (Table 4).

DISCUSSION

In Türkiye, the initial assessment of assault cases and the preparation of forensic reports are performed by emergency medicine physicians. Injuries can sometimes lead to dire consequences, including death. To reduce incidents of assault and their associated damages, the

Table 1. Distribution according to trauma location, symptoms, outcome, and ordered consultations.

	n	%
Trauma Localization		
Nasal	423	14.59
Maxilla	487	16.79
Mandible	242	8.34
Zygoma	217	7.48
Frontal	476	16.41
Orbit	417	14.38
Temporal	303	10.45
Parietal	279	9.62
Occipital	259	8.93
Thoracolumbar vertebrae	152	5.24
Cervical vertebrae	538	18.55
Thoracic	505	17.41
Abdomen	142	4.90
Pelvis	123	4.24
Upper Extremity	1391	47.97
Lower Extremity	632	21.79
Symptoms		
Vital Danger	23	0.79
Sensitivity	1811	62.45
Edema	662	22.83
Hematoma	147	5.07
Abrasion	1519	52.38
Ecchymosis	1093	37.69
Incision	590	20.34
Outcome		
Outpatient discharge	2625	90.52
Hospitalization in surgical departments	50	1.72
Outpatient clinic control	187	6.45
Treatment rejection	21	0.72
Unauthorized abandonment	17	0.59
Consultations		
Neurosurgery	38	1.31
General surgery	20	0.69
Obstetrics and gynecology	23	0.79
Eye diseases	107	3.69
Plastic surgery	58	2.00
ENT	241	8.31
Orthopedics	217	7.48
Thoracic surgery	9	0.31
CVS	9	0.31
Neurology	4	0.14
Urology	7	0.24
Social services	5	0.17
ENT: ears, nose, and throat, CVS: Cardiovascular Surgery		

WHO, non-governmental organizations, and governments have been compelled to implement specific measures¹².

In underdeveloped countries, there is an increase in the number of various assault incidents worldwide. In Türkiye, articles have been published on this issue. In a study conducted by Seviner et al.¹³, in the southern part of Türkiye, in which 5870 forensic cases were analyzed, it was determined that the number of male cases was approximately twice that of female cases. In a study conducted by Kanal et al.¹⁴ in another province of Türkiye, the sex ratios were strikingly similar to the findings of this study. Our study was conducted in western Türkiye (Istanbul), and the sex ratios were similar to those observed in these two studies. According to data from the Turkish Statistical Institute, the education level in Istanbul is higher than in the other two locations where the studies were conducted. It was considered particularly interesting that there was no difference in sex ratios between the studies conducted in the three regions with different education levels. This observation suggests that assault incidents are a general issue in Türkiye, regardless of educational level. Our study, conducted in one Istanbul region, which has been officially recorded as the area with the highest education level, aligns with this trend. Despite being in a region with a high education level, assault incidents did not show significant differences between areas with lower educational levels.

The average age of assault victims is a significant factor in cases of assault. In our study, the mean age of assault victims presenting to our emergency department was similar to the results of four studies¹⁴⁻¹⁷ conducted in different regions of Türkiye. These results indicate that assault incidents in Türkiye predominantly affect younger age groups.

In the study by Bennet et al.¹⁸, although the sex ratios were slightly different, they favor males. The study suggests that the reason for the higher involvement of males in assault incidents is their increased risk of violence. There are also hypotheses that the male-favoring ratios result from better interhemispheric connections in females and deficits in the frontal cortex in males¹⁹. Fazel et al.²⁰ also emphasized the association between assault incidents and psychiatric disorders. In our study, all 2900 male and female victims were assaulted by males. There were no cases in which the perpetrator was female. Although our research did not specifically investigate the causes of these incidents, we believe that the dominant aggression observed in males cannot be solely attributed to structural and functional differences in the brain. We consider that factors such as family structure, the upbringing of children, the content

Table 2. Distribution of trauma sites according to sex.

	All Patients n (%)	Male n (%)	Female n (%)	p+
Trauma localization				
Nasal	423 (14.59)	340 (18.38)	83 (7.90)	0.0001
Maxilla	487 (16.79)	326 (17.62)	161 (15.33)	0.113
Mandible	242 (8.34)	140 (7.57)	102 (9.71)	0.045
Zygoma	217 (7.48)	150 (8.11)	67 (6.38)	0.104
Frontal	476 (16.41)	350 (18.92)	126 (12.00)	0.0001
Orbit	417 (14.38)	298 (16.11)	119 (11.33)	0.0001
Temporal	303 (10.45)	211 (11.41)	92 (8.76)	0.025
Parietal	279 (9.62)	189 (10.22)	90 (8.57)	0.149
Occipital	259 (8.93)	173 (9.35)	86 (8.19)	0.292
Thoracolumbar vertebra	152 (5.25)	81 (4.39)	71 (6.77)	0.006
Cervical vertebrae	538 (18.55)	332 (17.95)	206 (19.62)	0.265
Thoracic	505 (17.41)	319 (17.24)	186 (17.71)	0.748
Abdomen	142 (4.90)	83 (4.49)	59 (5.62)	0.174
Pelvis	123 (4.24)	49 (2.65)	74 (7.05)	0.0001
Upper extremity	1.391 (47.97)	806 (43.57)	585 (55.71)	0.0001
Lower extremity	632 (21.79)	340 (18.38)	292 (27.81)	0.0001
Outcome				
Outpatient discharge	2625 (90.52)	1631 (88.16)	994 (94.67)	
Surgical departments	50 (1.72)	41 (2.22)	9 (0.86)	
Outpatient clinic control	187 (6.45)	150 (8.11)	37 (3.52)	
Treatment rejection	21 (0.72)	17 (0.92)	4 (0.38)	
Unauthorized abandonment	17 (0.59)	11 (0.59)	6 (0.57)	0.0001
Signs of injury				
Tenderness	1811(62.45)	1114(60.22)	697(66.38)	0.001
Edema	662(22.83)	466(25.19)	196(18.67)	0.0001
Hematoma	147(5.07)	109(5.89)	38(3.62)	0.007
Abrasion	1519(52.38)	1009(54.54)	510(48.57)	0.002
Ecchymosis	1093(37.69)	578(31.24)	515(49.05)	0.0001
Laceration	590(20.34)	471(25.46)	119(11.33)	0.0001
+:Chi-square test				

Table 3. Comparison of patient outcomes according to arrival time.

	All Hours	08:00-15:00	15:01-00:00	00:01-07:59	P
Age	35.53±12.46	37.53±12.50	35.42±12.66	33.56±11.54	0.0001*
Outcome					
Outpatient discharge	2625 (90.52)	650 (92.33)	1431 (91.15)	544 (86.90)	0.02+
Surgical departments	50 (1.72)	9 (1.28)	27 (1.72)	14 (2.24)	0.02+
Outpatient control	187 (6.45)	34 (4.83)	96 (6.11)	57 (9.11)	0.02+
Treatment rejection	21 (0.72)	4 (0.57)	11 (0.70)	6 (0.96)	0.02+
Vital danger	23 (0.79)	5 (0.71)	13 (0.83)	5 (0.80)	0.958+
*One-Way Analysis of Variance					

Table 4. Association between trauma locations and patient outcomes.						
	Discharge	Hospitalization in the surgical department	Outpatient control	Treatment rejection	Unauthorized abandonment	P-value
Nasal	325 (12.38)	5 (10.00)	84 (44.92)	7 (33.33)	2(11.76)	0.0001
Maxilla	432 (16.46)	14 (28.00)	33 (17.65)	7 (33.33)	1 (5.88)	0.035
Mandible	227 (8.65)	8 (16.00)	6 (3.21)	1 (4.76)	0 (0.00)	0.014
Zygoma	188 (7.16)	5 (10.00)	23 (12.30)	1 (4.76)	0 (0.00)	0.069
Frontal	432 (16.46)	11 (22.00)	23 (12.30)	5 (23.81)	5 (29.41)	0.173
Orbit	341 (12.99)	15 (30.00)	54 (28.88)	6 (28.57)	1 (5.88)	0.0001
Temporal	278 (10.59)	3 (6.00)	20 (10.70)	1 (4.76)	1 (5.88)	0.693
Parietal	256 (9.75)	4 (8.00)	16 (8.56)	2 (9.52)	1 (5.88)	0.949
Occipital	243 (9.26)	5 (10.00)	8 (4.28)	0 (0.00)	3 (17.65)	0.060
Toracolomber	133 (5.07)	5 (10.00)	13 (6.95)	1 (4.76)	0 (0.00)	0.343
Cervical	508 (19.35)	4 (8.00)	22 (11.76)	1 (4.76)	3 (17.65)	0.011
Thorax	460 (17.52)	7 (14.00)	31 (16.58)	4 (19.05)	3 (17.65)	0.968
Abdomen	121 (4.61)	5 (10.00)	12 (6.42)	3 (14.29)	1 (5.88)	0.085
Pelvics	110 (4.19)	3 (6.00)	9 (4.81)	1 (4.76)	0 (0.00)	0.859
Upper extremity	1.273 (48.50)	17 (34.00)	88 (47.06)	7 (33.33)	6 (35.29)	0.128
Lower extremity	563 (21.45)	24 (48.00)	37 (19.79)	4 (19.05)	4 (23.53)	0.0001

of the education provided, the social fabric of the society, the values and beliefs held, reward mechanisms, and the sanctity attributed to males contribute to these incidents.

The high prevalence of male violence is undoubtedly a global humanitarian issue that must be addressed. The fact that one in three women worldwide have been subjected to domestic violence by their partners confirms the severity of this problem.

As in the rest of the world, in Turkey, the years 2020-2021 were marked by a unique sociological atmosphere due to the COVID-19 pandemic and the measures taken, including quarantine applications. During this period, there was a decrease in hospital admissions because of the risk of COVID-19 transmission, and hospital operations changed^{21,22}. In our study, the lowest number of forensic case reports occurred in 2020. We attribute this to the strict implementation of the COVID-19 pandemic restrictions in our country during that year. A similar result was demonstrated in a study by Kuitunen and colleagues²³. This is likely due to the first lockdown implemented in Turkey on April 10-12, 2020. In the following two years, an increase in cases was observed. Faez et al. also noted that lockdowns and curfews decreased hospital admissions for forensic cases²⁴. However, it has become controversial whether restrictions on social life increase domestic violence incidents.

It is well-known that the number of trauma admissions to the emergency department increases during the summer months and prolonged vacations²⁵. In our study, when the seasonal distribution of assault cases was analyzed, it was observed that approximately one-third of all cases occurred during the summer months. Similar findings have been reported by Küçüker et al.²⁶, Altun et al.²⁷ also reported cases in different regions of Türkiye, indicating that forensic cases mainly occur during summer. The increase in forensic cases during the summer months in Türkiye can be attributed to the intensified workload in specific service sectors, particularly agriculture and livestock, during this period. Examining the hours of forensic case admissions have been a subject of interest in both our study and many others. In a study by Korkmaz et al.²⁸, who screened 47.758 cases, forensic case admissions mainly occurred during afternoon and night shifts. We conclude that factors such as fatigue at the end of the day, heavy traffic during rush hours in large cities leading to stress, higher alcohol consumption and its consequences during late-night hours, conflicts among fans attending late-night sports events, and relatively lower tolerance limits among young partners contribute to triggering assault incidents.

In cases of assault, the localization of injuries is crucial for assessing the victim's risk of mortality, organ damage/function loss, intention of the assailant and the degree of punishment they may face. Therefore, the first responding

physician must handle the situation exceptionally sensitively and maintain accurate records. In a study by Brink et al.²⁹, which analyzed 1481 cases, more than 95% of the victims were exposed to blunt trauma, and the trauma locations occurred in the craniofacial region in 69% of cases. In Payne-James and Dean's study, craniofacial injuries were identified in at least half of the patients³⁰. Our study confirms the literature findings of a high incidence of craniofacial injuries, especially in the nasal, frontal, orbital, and maxillary regions. In attacks involving face-to-face violence, the craniofacial region is often the primary target. The higher incidence of nasal, maxillary, and orbital trauma in male patients than in female patients can be explained in this context. Conversely, women are more likely to find themselves in defensive positions, which may influence the localization of the impact they receive. Spinal injuries have been observed more frequently in women than in men. In our study, statistically significantly higher rates of spinal injuries were observed in female patients than in male patients. In a prospective study conducted in many different centers in Türkiye, examining approximately 150,000 cases, the most frequently identified injury regions were extremities, head-neck region, and abdomen³¹. We believe that the high frequency of extremity injuries, second only to head traumas, is mainly due to the use of extremities as both defensive and offensive tools.

It is well known that the mortality rate of male assault cases is higher. Data for approximately 25 years of assault cases in Scotland revealed that mortality is higher in men than in women³², no fatal outcomes were observed in our study. However, the proportion of male patients was significantly higher among those in whom life-threatening conditions were detected upon admission and hospitalization to surgical departments. The higher rate of outpatient discharge for female patients compared with male patients also indicates that males tend to sustain more serious injuries during assaults. Consistently, more severe injuries, such as cuts and fractures, were observed in male patients than in female patients.

Generally, nine out of every ten patients in our study were directly discharged from the hospital in a multicenter study conducted by Seviner et al.¹³, which involved forensic cases presenting to the emergency department; 56.8% of the cases were hospitalized, and 21.1% were observed to be in a life-threatening condition. In our study, the rate of life-threatening conditions was lower than that in this study (0.71%). A study in Brazil involving 62 centers and 4835 cases emphasized that

young males caused more severe and fatal injuries using cutting or stabbing tools and firearms³³. Some studies demonstrated the influence of sociocultural factors, societal patriarchal norms, and gender roles, suggesting that violence among men can lead to more dangerous consequences³⁴.

Since the injury locations are generally concentrated in the head and extremities, the need for consultation is usually in the otorhinolaryngology departments, plastic surgery, neurosurgery, and orthopedics. It is important to note that the mean age of patients consulted and hospitalized in surgical departments was higher than that of patients discharged as out-patients. There could be many reasons for this that we could not identify. One of the reasons may be that body tissues become more vulnerable with age, and accordingly, more injuries are sustained during impacts.

Limitations

Our study was designed retrospectively, and detailed data regarding the mechanisms of trauma formation could not be collected. The relationship between trauma and specific subgroups, such as the elderly population's abuse or the association with subgroups like partner violence, has not been investigated. Only adult patients (>16 years of age) were included, and incidents of abuse and violence in children were not analyzed. The primary cause of the assault could not be determined. This study is not multicenter and focuses on only one region of Istanbul, which is a metropolitan city.

CONCLUSION

In our study, the following findings were obtained: All assault incidents were committed by male individuals, with males being more frequently and severely assaulted than females. The victims were predominantly from the youth group. Assault incidents primarily occurred during the summer months and outside working hours. The head-neck region and extremities were the most injured areas during assaults. Assault incidents have increased over the years. Except for mortality, our findings demonstrated similarities with the literature. No fatalities were observed, and nine out of ten patients admitted to the emergency department were discharged home.

Ethics

Ethics Committee Approval: This study was approved by the H.S. Istanbul Medeniyet University Göztepe Training and Research Hospital Clinical Research Ethics Committee (decision no :2023/0219, date: 29.03.2023).

Informed Consent: Presented to the emergency department of tertiary hospital with a "history of assault" were retrospectively examined between January 01, 2020 and December 31, 2022

Footnote

Author Contributions

Concept: M.Y.S., B.A., E.T., Design: M.Y.S., B.A., G.A.S., Data Collection and/or Processing: M.Y.S., B.A., V.C., E.T., Analysis and/or Interpretation: M.Y.S., K.A., B. A.C., N.S., Literature Search: K.A., B.A.C., N.S., Writing: M.Y.S., B.A., G.A.S.

Conflict of Interest: The authors have no conflict of interest to declare.

Financial Disclosure: The authors declared that this study has received no financial support.

REFERENCES

1. Zufer I, Fix RL, Stone E, et al. Documentation of Trauma-Informed Care Elements for Young People Hospitalized After Assault Trauma. *J Surg Res.* 2024;296:665-73.
2. Mercy JA, Hillis SD, Butchart A, et al. Interpersonal Violence: Global Impact and Paths to Prevention. In: Mock CN, Nugent R, Kobusingye O, Smith KR, eds. *Injury Prevention and Environmental Health.* 3rd ed. Washington (DC): The International Bank for Reconstruction and Development / The World Bank; October 27.
3. Procaccia R, Castiglioni M. The mediating effect of cognitive and emotional processing on PTSD and depression symptoms reduction in women victims of IPV. *Front Psychol.* 2022;13:1071477.
4. Wankhede AG. Patterned injuries caused by a wooden plank. *J Forensic Leg Med.* 2008;15:118-23.
5. tochero L, Pinto LW. Prevalence of violence against women living in rural areas and associated factors: a cross-sectional study based on the 2019 National Health Survey. *Cien Saude Colet.* 2024;29:20452022.
6. Matzopoulos R, Bowman B, Butchart A, et al. The impact of violence on health in low- to middle-income countries. *Int J Inj Contr Saf Promot.* 2008;15:177-87.
7. İlçe A, Erkol MH, Alpteker H, et al. Retrospective Analysis of Forensic Case Reports Who Had Applied to the Emergency Service in the City Centre of Bolu. *Abant Med J.* 2018;7:68-75.
8. Bardaa S, Dhoub H, Karray N, et al. Intentional Interpersonal Violence: Epidemiological and analytical study about 973 cases at the forensic unit in Habib BOURGUIBA University Hospital in Sfax, Tunisia. *Forensic Science International: Reports.* 2020;2:100106.
9. Kafadar H. The Evaluation of Assaults With Social and Medico-legal Aspect in 3172 Cases. *Mitt Klosterneuburg Rebe Wein Obstb Fruchteverwert.* 2014;64:100-5.
10. Alabdulqader S, Alabdulqader R, Madadin M, et al. Emergency Physicians' Awareness of Medico-Legal Case Management: A Cross-Sectional Study from Saudi Arabia. *Saudi J Med Med Sci.* 2023;11:60-6.
11. Teyin A, Kaya A, Şenol E, et al. Our opinion about the guide entitled "Evaluation of the wounding crimes described by Turkish Criminal Code with Forensic Science Perspective." *Anatol J Med.* 2014;24:219-21.
12. Sugg N. Intimate partner violence: prevalence, health consequences, and intervention. *Med Clin North Am.* 2015;99:629-49.
13. Seviner M, Kozaci N, Ay MO, et al. Analysis of Judicial Cases at Emergency Department. *Cukurova Medical Journal.* 2013;38:250-60.
14. Kanal, T, Türkmen İnanır, N, Özdemir, et al. Evaluation of The Forensic Cases Admitted to the Emergency Department of Bursa Uludag University Medical Faculty in 2020. 4th International 20th National Forensic Sciences Congress (pp.258). İzmir, Turkey; 2023.
15. Duman A, Kapçı M, Bacakoğlu G, et al. Evaluation of trauma patients in emergency department. *Med J SDU.* 2014;21:45-8.
16. Serinken M, Türkçüler İ, Aydın B, et al. Application times of the forensic cases to the emergency department. *Eurasian J Emerg Med.* 2010;9:89-92.
17. Hakkoymaz H, Keten HS, Artuç S, et al. Evaluation of Medico-Legal Reports in Respect of the Turkish Penal Code. *J Kartal TR.* 2014;25:177-80.
18. Bennett S, Farrington D, Huesmann L. Explaining Gender Differences in Crime and Violence: The Importance of Social Cognitive Skills. *Aggress Violent Behav.* 2005;10:263-88.
19. Takahashi A, Nagayasu K, Nishitani N, et al. Control of Intermale Aggression by Medial Prefrontal Cortex Activation in the Mouse. *PLOS ONE.* 2014;9:94657.
20. Fazel S, Wolf A, Chang Z, et al. Depression, and violence: a Swedish population study. *Lancet Psychiatry.* 2015;2:224-32.
21. Giostra F, Mirarchi MG, Farina G, et al. Impact of COVID-19 pandemic and lockdown on emergency room access in Northern and Central Italy. *Emerg Care J.* 2021;17.
22. Ayşenur DB, Tuğba D, İlhan İ. Normative Analysis and Ethical Evaluation of Curfew and Restrictions as a Public Health Measure During the Covid-19 Pandemic. *JARHS.* 2021;4:1-18.
23. Kuitunen I, Ponkilainen VT, Launonen AP, et al. The effect of national lockdown due to COVID-19 on emergency department visits. *Scand J Trauma Resusc Emerg Med.* 2020;28:114.
24. Díaz-Faes DA, Vidal-Codina F, Segura A, et al. How the COVID-19 pandemic hit crime in Barcelona: Analysis of variation in crime trends. *Eur J Criminol.* 2023;20:792-816.
25. Dagar S, Sahin S, Yilmaz Y, et al. Emergency Department During Long Public Holidays. *Turk J Emerg Med.* 2014;14:165-71.
26. Evaluation Of Traffic Accident Cases Admitted To The Emergency Department Of The Firat University Hospital In 1997-2001. *Turk J Emerg Med.* 2003;3:11-15.
27. Altun G, Azmak AD, Yılmaz A, et al. The characteristics of the cases admitted to the Emergency Department of Trakya University Medical Faculty. *Adli Tıp Bülteni.* 1997;2:62-6.
28. Korkmaz T, Kahramansoy N, Erkol Z, et al. Evaluation of the Forensic Patients Presenting to the Emergency Department and Legal Reports. *Medical Bulletin of Haseki.* 2012;50:14-20.

29. Brink O, Vesterby A, Jensen J. Pattern of injuries due to interpersonal violence. *Injury*. 1998;29:705-9.
30. Payne-James JJ, Dean PJ. Assault and injury in clinical forensic medical practice. *Med Sci Law*. 1994;34:202-6.
31. Acin M, Karakisa H, Satar S. Demographic Analysis and Intentionality of Injuries in the Emergency. *Phnx Med J*. 2020;2:152-9.
32. Leyland AH, Dundas R. The social patterning of deaths due to assault in Scotland, 1980–2005: population-based study. *J Epidemiol Community Health*. 2010;64:432-9.
33. Gawryszewski V, Silva M, Carvalho Malta D, et al. Violence-related injury in emergency departments in Brazil. *Rev Panam Salud Publica*. 2008;24:400-8.
34. Al Kiyumi MH, Al Shidhani AS, Al Sumri H, et al. Intimate Partner Violence in Khaliji Women: A Review of the Frequency and Related Factors. *Int J Environ Res Public Health*. 2023;20:6241.