

AUTHOR'S REPLY

To the Editor,

We would like to express our gratitude to the researcher who took the time to read our article titled “The relationship between serum vitamin D level and asthma” and added value to our writing by writing to the editor. Contacted by a valuable researcher, “Patients with inhaled steroids may have decreased serum vitamin D” information is available in the literature [1, 2]. In this respect, the researcher is extremely right. Thus, we included patients with newly diagnosed and not receiving treatment (inhaled steroids and others). We mentioned this in the study design section of the material and method section [3].

 Mehmet Yasar Ozkars

Department of Pediatric Immunology and Allergy,
Gaziantep University Faculty of Medicine,
Gaziantep, Turkey

E-mail: myozkars@hotmail.com

doi: 10.14744/nci.2020.88964



REFERENCES

1. Searing DA, Zhang Y, Murphy JR, Hauk PJ, Goleva E, Leung DY. Decreased serum vitamin D levels in children with asthma are associated with increased corticosteroid use. *J Allergy Clin Immunol* 2010;125:995–1000. [\[CrossRef\]](#)
2. Stelmach I, Olszowiec-Chlebna M, Jerzynska J, Grzelewski T, Stelmach W, Majak P. Inhaled corticosteroids may have a beneficial effect on bone metabolism in newly diagnosed asthmatic children. *Pulm Pharmacol Ther* 2011;24(4):414–20. [\[CrossRef\]](#)
3. Ozkars MY, Keskin O, Almacioglu M, Kucukosmanoglu E, Keskin M, Balci O. The relationship between serum vitamin D level and asthma. *North Clin Istanbul* 2019;6:334–40. [\[CrossRef\]](#)