

Supracondylar extra-articular femur fracture after cementless unicompartmental knee replacement: A rare complication

 Doğaç Karagüven, M.D.,¹  Tuğrul Yıldırım, M.D.,²  Burak Akan, M.D.,¹
 Mahmut Nedim Doral, M.D.¹

¹Department of Ortopedics and Traumatology, Ufuk University Faculty of Medicine, Ankara-Türkiye

²Department of Ortopedics and Traumatology, Hand Microsurgery Orthopedics Traumatology Hospital, İzmir-Türkiye

ABSTRACT

This case report defines an infrequent complication of unicompartmental knee replacement. Periprosthetic supracondylar femoral fracture after total knee replacement is a challenging problem for orthopedic surgeon. To the best of our knowledge, this is the only case describing periprosthetic supracondylar femoral fracture after unicompartmental knee replacement.

Keywords: Complication; periprosthetic fracture; supracondylar femoral fracture; unicompartmental knee replacement.

INTRODUCTION

Oxford unicompartmental knee replacement (UKR) has become an alternative procedure for osteoarthritis of the knee with excellent long-term results.^[1,2] UKR is a safe procedure, and low perioperative complication rates have been reported in the literature.^[3] The most common complications are aseptic loosening, polyethylene dislocations, unexplained pain, and periprosthetic tibial fractures.^[2,4] The prevalence of supracondylar fractures after total knee replacement (TKR) is reported to be 0.5–2%.^[5,6] Although few case reports have been published on condylar fractures; to the best of our knowledge, periprosthetic supracondylar femoral fracture after UKR has not been reported in the literature.^[7–9]

In this case report, we present a patient with a post-operative supracondylar femoral fracture who had been previously treated for ipsilateral condylar periprosthetic femur fracture after UKR. We aimed to discuss the mechanism of the fracture, treatment strategies, and its impact on patient's clinical outcome (A written informed consent was obtained from the patient for publication of this case report).

CASE REPORT

A 53-year-old woman underwent cementless mobile-bearing Oxford partial knee phase 3 (Biomet Orthopedics) by the senior author in 2012 for anteromedial osteoarthritis. One year after an uneventful post-operative recovery, the patient fell on the ground level, while walking on the street. A minimally displaced medial femoral condyle fracture was diagnosed without ligamentous instability, and a closed reduction and percutaneous fixation with 6.5 mm cannulated cancellous lag screws were performed.^[7] Before the fracture, the patient's oxford knee score (OKS) was 46. After the fracture healing, her OKS was 42. In 2014, the patient had a second fall at ground level, which resulted in the unified classification system type C supracondylar femoral fracture^[10] (Fig. 1). Both femoral and tibial components seemed well-fixed, and insert dislocation was not observed in X-rays. We performed an open reduction and internal fixation and used compression screws and anatomical distal femoral locking plate for fixation (Fig. 2). Active and passive range-of-motion exercises were initiated immediately after the surgery. Union was observed at the fracture site at 12 weeks on X-rays, and full weight-

Cite this article as: Karagüven D, Yıldırım T, Akan B, Doral MN. Supracondylar extra-articular femur fracture after cementless unicompartmental knee replacement: A rare complication. *Ulus Travma Acil Cerrahi Derg* 2022;28:1754-1757.

Address for correspondence: Mahmut Nedim Doral, M.D.

Ufuk Üniversitesi Tıp Fakültesi, Ortopedi ve Travmatoloji Anabilim Dalı, Türkiye

Tel: +90 312 - 586 70 00 E-mail: mndoral@gmail.com

Ulus Travma Acil Cerrahi Derg 2022;28(12):1754-1757 DOI: 10.14744/tjtes.2021.03435 Submitted: 07.03.2021 Revised: 09.03.2021 Accepted: 01.05.2021
 OPEN ACCESS This is an open access article under the CC BY-NC license (<http://creativecommons.org/licenses/by-nc/4.0/>).






Figure 1. Type C supracondylar femoral fracture.




Figure 2. After open reduction and internal fixation with compression screws and anatomical distal femoral locking plate.




Figure 3. Four years after surgery, the implants were still well-fixed.

bearing was allowed. The patient had 130° range of motion and was able to walk independently 6 months after surgery. Four years after surgery, the implants were still well-fixed, and OKS was 36 (Fig. 3).

DISCUSSION

This case report proves unicompartmental prosthesis that remains intact even after two subsequent periprosthetic fractures,

and fracture healing can be achieved without complications. The clinical outcome was excellent and the patient regained almost full range of motion. To the best of our knowledge, no cases have been reported in the literature regarding such a fracture.

The goal of managing these injuries includes restoration of axial alignment and length and stability to allow early mobilization. However, achieving this goal remains to be challenging for orthopedic surgeons. Due to poor bone quality and fracture location, non-union is not uncommon with periprosthetic distal femoral fractures after TKR, and non-union rates have been reported between 0% and 50%.^[11] Preservation of soft tissue and osseous vascularity has been recommended as much as possible to reduce non-union rates.^[12] We think that the same principles should be applied in the surgical management of periprosthetic fractures after UKR. Restoring axial alignment was especially crucial in our case, because we think that unicondylar prosthesis would less tolerate shear forces than TKR caused by malalignment.

Fracture localization and displacement are essential considerations in the management of supracondylar periprosthetic femur fractures. Despite the fact that, the use of locked plating and retrograde intramedullary nailing (IMN) has been recommended for displaced supracondylar femoral fractures after TKR, no consensus exists regarding the ideal treatment strategy. We think that both strategies are applicable for periprosthetic femoral fractures after UKR. Retrograde IMN offers more stability in the presence of comminution of the medial cortex but is limited by poor cancellous bone, type of prosthesis, and pre-existing hardware in the proximal femur.^[13,14] Due to the inherent advantage of UKR design that the femoral notch is not occupied by the femoral component, IMN may be more applicable. Locked plating has been recommended for low supracondylar fractures due to difficulties in achieving distal fixation with IMN and controlling varus collapse.^[15] In a recent meta-analysis, Li et al.^[16] compared the clinical results of locked plates and retrograde IMN and found no statistically significant difference in terms of union, operating time, and rates of complication.

Several factors can be attributed to the fractures in our patient. The replacement of an intramedullary rod during the procedure may have created a stress riser effect, which eventually diminished the structural strength of the femur.^[7] Subsequent fractures with minor trauma may indicate that osteoporosis might have played a role in the etiology. However, our patient did not have any known patient-related risk factors or previous fractures related to osteoporosis.

Meniscal bearing dislocation is a well-known complication of UKR with mobile bearing insert due to inappropriate ligament balance and component replacement.^[16] Despite the fact that the patient's knee was exposed to shear forces during trauma, the insert remained in its place. We think mobile-

bearing causes more compressive and less tensile and shear force on the implants, thus avoiding an excessive load at the bone-implant interfaces. Thanks to this mechanism, components may not be loose despite the load transferred during the trauma.

Although many systems are available for classifying periprosthetic fractures after TKR, there is no such classification regarding UKR periprosthetic fractures.^[17] The reason for not needing such a classification may be due to the relatively low incidence of periprosthetic fractures after UKR than TKR. However, UCR can be considered as the most applicable and inclusive classification for periprosthetic fracture after UKR.^[10] This system is based on the Vancouver classification and has been defined as a relatively simple alternative that can be used to describe any periprosthetic fracture.^[10,18] According to UCR classification, our case had Type C fracture, in which the fracture line was distant to the bed of the implant. It was suggested that these fractures could be managed open or closed osteosynthesis without involving the implant as we did in our case.^[19]

Conclusion

Despite subsequent fractures, if UKR is properly replaced and has appropriate ligament balance, insert dislocation would not occur, and the components would remain intact. In addition, bone healing could be achieved with an excellent clinical outcome. The treatment goal should be the restoration of alignment and achieving stability to allow early mobilization.

Informed Consent: Written informed consent was obtained from the patient for the publication of the case report and the accompanying images.

Peer-review: Externally peer-reviewed.

Authorship Contributions: Concept: D.K., T.Y., B.A., M.N.D.; Design: D.K., T.Y., B.A., M.N.D.; Supervision: D.K., T.Y., B.A., M.N.D.; Data: D.K., B.A.; Analysis: D.K., B.A., M.N.D.; Literature search: D.K., T.Y.; Writing: D.K., T.Y.; Critical revision: D.K., T.Y., B.A., M.N.D.

Conflict of Interest: None declared.

Financial Disclosure: The authors declared that this study has received no financial support.

REFERENCES

- Pandit H, Jenkins C, Gill HS, Barker K, Dodd CA, Murray DW. Minimally invasive Oxford phase 3 unicompartmental knee replacement: Results of 1000 cases. *J Bone Joint Surg Br* 2011;93:198–204. [\[CrossRef\]](#)
- Price AJ, Waite JC, Svard U. Long-term clinical results of the medial

- Oxford unicompartmental knee arthroplasty. *Clin Orthop Relat Res* 2005;435:171–80. [\[CrossRef\]](#)
- Morris MJ, Molli RG, Berend KR, Lombardi AV Jr. Mortality and perioperative complications after unicompartmental knee arthroplasty. *Knee* 2013;20:218–20. [\[CrossRef\]](#)
- Vince KG, Cyran LT. Unicompartmental knee arthroplasty. New indications, more complications? *J Arthroplasty* 2004;19:9–16. [\[CrossRef\]](#)
- Ayers DC, Dennis DA, Johanson NA, Pellegrini JV. Instructional course lectures, The American academy of orthopaedic surgeons-common complications of total knee arthroplasty. *J Bone Joint Surg Am* 1997;79:278–311. [\[CrossRef\]](#)
- Meek RM, Norwood T, Smith R, Brenkel IJ, Howie CR. The risk of periprosthetic fracture after primary and revision total hip and knee replacement. *J Bone Joint Surg Br* 2011;93:96–101. [\[CrossRef\]](#)
- Akan B, Yildirim T, Karagüven D. Medial femoral condyle fracture after cementless unicompartmental knee replacement: A rare complication. *Knee* 2013;20:295–97. [\[CrossRef\]](#)
- Kim KT, Lee S, Cho KH, Kim KS. Fracture of the medial femoral condyle after unicompartmental knee arthroplasty. *J Arthroplasty* 2009;24:1143.e21–4. [\[CrossRef\]](#)
- Ten Brinke B, De Haan LJ, Koenraadt KL, Van Geenen RC. Medial femoral condyle fracture as an intraoperative complication of Oxford unicompartmental knee replacement. *Knee Surg Sports Traumatol Arthrosc* 2016;24:3191–93. [\[CrossRef\]](#)
- Duncan CP, Haddad FS. The unified classification system (UCS): Improving our understanding of periprosthetic fractures. *Bone Joint J* 2014;96–B:713–6. [\[CrossRef\]](#)
- Hoffmann MF, Jones CB, Sietsema DL, Koenig SJ, Tornetta P. Outcome of periprosthetic distal femoral fractures following knee arthroplasty. *Injury* 2012;43:1084–9. [\[CrossRef\]](#)
- Bolhofner BR, Carmen B, Clifford P. The results of open reduction and internal fixation of distal femur fractures using a biologic (indirect) reduction technique. *J Orthop Trauma* 1996;10:372–7. [\[CrossRef\]](#)
- Herrera DA, Kregor PJ, Cole PA, Levy BA, Jönsson A, Zlowodzki M. Treatment of acute distal femur fractures above a total knee arthroplasty: Systematic review of 415 cases (1981-2006). *Acta Orthop* 2008;79:22–7. [\[CrossRef\]](#)
- Chettiar K, Jackson MP, Brewin J, Dass D, Butler-Manuel PA. Supracondylar periprosthetic femoral fractures following total knee arthroplasty: Treatment with a retrograde intramedullary nail. *Int Orthop* 2009;33:981–5. [\[CrossRef\]](#)
- Matlovich NF, Lanting BA, Vasarhelyi EM, Naudie DD, McCalden RW, Howard JL. Outcomes of surgical management of supracondylar periprosthetic femur fractures. *J Arthroplasty* 2017;32:189–92. [\[CrossRef\]](#)
- Li B, Gao P, Qiu G, Li T. Locked plate versus retrograde intramedullary nail for periprosthetic femur fractures above total knee arthroplasty: A meta-analysis. *Int Orthop* 2016;40:1689–95. [\[CrossRef\]](#)
- Jeong JH, Kang H, Ha YC, Jang EC. Incarceration of a dislocated mobile bearing to the popliteal fossa after unicompartmental knee arthroplasty. *J Arthroplasty* 2012;27:323.e5–7. [\[CrossRef\]](#)
- Rorabeck CH, Taylor JW. Classification of periprosthetic fractures complicating total knee arthroplasty. *Orthop Clin North Am* 1999;30:209–14. [\[CrossRef\]](#)
- Konan S, Sandiford N, Unno F, Masri BS, Garbuz DS, Duncan CP. Periprosthetic fractures associated with total knee arthroplasty: An update. *Bone Joint J* 2016;98–B:1489–96. [\[CrossRef\]](#)

OLGU SUNUMU - ÖZ

Çimentosuz unikondiler diz protezi sonrası suprakondiler femur kırığı: Nadir bir komplikasyon

Dr. Doğaç Karagüven,¹ Dr. Tuğrul Yıldırım,² Dr. Burak Akan,¹ Dr. Mahmut Nedim Doral¹

¹Ufuk Üniversitesi Tıp Fakültesi, Ortopedi ve Travmatoloji Anabilim Dalı, Ankara
²EMOT Hastanesi, Ortopedi ve Travmatoloji Kliniği, İzmir

Bu olgu sunumunda unikondiler diz protezinin seyrek görülen bir komplikasyonu tanımlanmaktadır. Total diz protezi cerrahisi sonrası oluşabilen periprotetik suprakondiler femur kırıkları, ortopedi cerrahları için başedilmesi zor bir komplikasyondur. Bildiğimiz kadarıyla bu olgu sunumu, unikondiler diz protezi sonrası gerçekleşen tek periprotetik suprakondiler femur kırığıdır.

Anahtar sözcükler: Komplikasyon; periprotetik kırık; suprakondiler femur kırığı; unikondiler diz protezi.

Ulus Travma Acil Cerrahi Derg 2022;28(12):1754-1757 doi: 10.14744/tjtes.2021.03435