

- 4) NF Sethna, D Zurakowski , RM Brostowicz, J Bacsik, LJ Sullivan, F Shapiro. Tranexamic acid reduced intraoperative blood loss in pediatric patients undergoing scoliosis surgery. *Anaesthesiology* 2005;102:727-732.
- 5) Ian Crabb. Anaesthesia for spinal surgery. *Anaesthesia and intensive care medicine*.- 2008; 10-1: 18-21.
- 6) EG Dawson, JE Sherman, LEA Kanim, MR Nuwer. Spinal cord monitoring: results of the Scoliosis Research Society and the European Spinal Deformity Society Survey. *Spine* 1991;16:5361-5364.
- 7) J Wong, H El Beheiry, YR Rampersaud, et al : Tranexamic acid reduces perioperative blood loss in adult patients having spinal fusion surgery. *Anesth Analg* 2008; 107(5),1479-1486.
- 8) A Sharma, JC Lawmin, MG Irwin. Anaesthesia for spinal surgery. *Orthopaedic Anaesthesia*. 2012;13-3: 99-101.
- 9) Y Gürkan, A Eroglu, E Kelsaka, H Kursad, A Yılmazlar. Anaesthesia for Scoliosis Surgery. *Turk J. Anaesth Reanim.* 2013;41:88-97.
- 10) JF Cassady, G Lederhaas, et al. A randomized comparison of the effects of continuous thoracic epidural analgesia and intravenous patient controlled analgesia after posterior spinal fusion in adolescents. *Reg Anaesth Pain Med.* 2000;25:246-253.
- 11) M Szabo, W Denman, J Morota, J Roberts . Evaluation of airway edema in patients operated on in the prone position. *J Neurosurg Anaesthesiol* 1997;9:380.
- 12) HC Sagi, W Beutler, E Carroll, PJ Connolly. Airway complications associated with surgery on the anterior cervical spine. *Spine* 2002;27: 949-953.
- 13) Y Miura ,K Mimatsu , H Iwata . Massive tongue swellingas a complication after spinal surgery. *J Spinal Disord* 1996;9:339-341.